

A review of number lines, interval notation, and set notation

This is a general review of these concepts. We will look at how to represent solutions for problems like:

$$x > 5 \quad \text{or} \quad -3 \leq x < 4 \quad \text{or} \quad x > 2 \text{ or } x < -3$$

Use the example $x < 4$ to describe the 3 types of inequality notation. Write down any special symbols associated with it.

Number line:

The special signs/symbols are:

Graph $x < 4$ on a number line:

Interval Notation:

The special signs/symbols are:

Write $x < 4$ in interval notation:

Set Notation

The format of this is always:

Write $x < 4$ in set notation:

Examples Write the following inequalities on a number line, in interval notation, and in set notation.

a. $x \geq 5$

b. $x < -2.5$

c. $-4 \leq x < 7$

d. $x > 4$ or $x < -5$

Examples Write the following inequalities on a number line, in interval notation, and in set notation.

a. $x \leq -3$

b. $x > 12$

c. $0 < x \leq 8$

d. $x > 3$ or $x < 1$

Lesson: Exponents and the Order of Operations

Exponents

Label the parts of this exponent:

$$2^3$$

What does 2^3 literally mean?

Example 1

Evaluate the following exponents

a. 3^3

b. 5^3

c. 2^4

We need to understand exponents so that we can fully understand the order of operations. The order of operations is one of the handiest concepts in arithmetic that is often confused.

Order of Operations

Write down what the order of operations is:

Classic Example of the Order of Operations

Evaluate: $10 \div 5 \times 6$

Example 2

Evaluate the following. This page and the next page are both blank so that you have plenty of space to work out your calculations by hand.

a. $3 + 2^3 - 6 \div 3 \times 7$

b. $2 + 3^2 + 4 \times 2 - 5 \times 2$

c. $3(1 + 2^2 - 3)^2$

Examples: The Order of Operations

a. $10 - 3(1 + 2 \times 3) - 2^2 \times 4$

b. $3(1 + 2^2 - 3)^2 + 5(4 - 2 \times 3)$

c. $\frac{1}{2}(5 + 3) - 2^2(5 - 6)$

Lesson: Advanced Notes on Exponents

Multiplicative Property of Negative One

Explain what is the difference between -5^2 and $(-5)^2$ from the standpoint of the order of operations:

Example 1

Evaluate the following.

a. $(-3)^2$

d. -3^3

b. $(-3)^3$

e. -7^2

c. -3^2

f. $(-2)^5$

Example 2

What is the sign on the following problems? Do not worry about the final answer, you only need to circle if the final answer is positive or negative.

a. $(-4)^7$ Positive Negative

d. $(-13)^{72}$ Positive Negative

b. $(-58)^{12}$ Positive Negative

e. -64^{14} Positive Negative

c. -72^{57} Positive Negative

f. $(-23)^{91}$ Positive Negative

There is one final rule we need to discuss.

Zero Exponents

For any real number a ,

Example 3

Evaluate the following:

a. 5^0

b. x^0

c. 35^0

d. -6^0

e. $(-7)^0$

f. $3^0 + 7^0$

g. $-5^0 + 2^0$

h. $(-3 + \sqrt{52})^0$

Lesson: Algebraic Expressions

First we will define the terminology, which is already written out for you.

- **Constant:** any real number in an expression that is not attached to a variable
- **Variable:** a letter that can stand for any set of real numbers, and we can substitute values in for it if desired
- **Coefficient:** a number that is being multiplied by a variable
- **Term:** this is a single number, variable, or a number and a variable multiplied together, often separated by a plus + or minus – sign.
- **Factor:** a number or variable that is being multiplied with another number or variable
- **Algebraic Expression:** a mathematical string of constants and terms that can be added, subtracted, multiplied or divided together

This terminology will be used often, so it's important to understand what it means.

Example 1

Classify each part of the expressions

<u>Expression</u>	<u>Constants</u>	<u>Variables</u>	<u>Coefficients</u>	<u>Terms</u>	<u>Factors</u>
a. $5x$					
b. $2x + y$					
c. 8					
d. x					
e. $2x^2 - 4x + 7$					

Algebraic expressions are often used in everyday situations; we just don't realize it.

Example 2

You want to go to the movies, and on Tuesdays a theatre runs a deal so that all tickets cost \$8. How much would it cost for each of the following situations?

- | | |
|--|--|
| a. A ticket for only you? | c. Tickets for 5 people, a giant tub of popcorn that costs \$20. |
| b. A ticket for you and 2 of your friends? | d. Tickets for z people? |

In the last part of this example, we wrote an algebraic expression where z represented some quantity. Notice that if you input any of the previous number of tickets from examples a, b, and c into z , you would get the same answers.

To evaluate expression, you can simply plug in the desired values for a variable.

Example 3

Evaluate each expression for $x = -2$, $y = 4$, and $z = \frac{1}{3}$ as applicable:

a. $3x - 7y$

b. $6z - x^2 + 2y$

c. $x + z^2$

Lesson: The Commutative, Associative, and Distributive Properties

Your math homework and English homework are due on the same day. Does it matter which one you do first?

This situation illustrates the idea of the Commutative Law, which we can now state below:

Commutative Law

Let a and b represent any two real numbers. Now we can define the following properties:

- The Commutative Law for Addition:

- The Commutative Law for Multiplication:

Example 1

Rewrite the following using the commutative law.

a. $3 + y$

b. $3c$

c. $\frac{2}{5}k$

d. $2 - z$

You are buying chips, a drink, and a notebook. Does it matter what order the cashier rings up this items in?

These situations illustrate the idea of the **Associative Law**, which we can now state below:

Let a , b , and c represent any three real numbers. Now we can define the following properties:

- The Associative Law for Addition:

- The Associative Law for Multiplication:

Example 2

Rewrite the following at least 2 different ways using the associative law.

a. $4xz$

b. $8 + 2 + 7 + 3$

These rules are helpful when we are computing longer expressions.

How to Simplify Calculations Using the Commutative and Associative Laws

- a. Suppose we want to calculate: $2 + 3 + 8 + 7$

Using the commutative law, we more cleverly group these numbers together like this:

$$2 + 8 + 3 + 7$$

Usually we recognize 10s faster than some other quantities, so from here we can rearrange and group like this to find the final answer:

- b. How can you rearrange $\left(\frac{1}{5} * \frac{1}{3}\right) \times (10 * 6)$ to make the calculations simpler to work with?

The Distributive Law

Let a , b , and c represent any three real numbers. Then the following property holds:

The distributive property effectively gives us permission to either add or multiply certain expressions in the order that simplifies the calculation.

Example 3

Use the distributive property as needed to simplify the following calculations.

a. $10\left(\frac{2}{5} - \frac{1}{2}\right)$

b. $\frac{2}{3}(6 + 9)$

c. $20\left(\frac{1}{4} + \frac{2}{5}\right)$

d. $\frac{2}{5}(14 - 4)$

The distributive law works in the opposite direction as well. If we start with $ab + ac$, according to the distributive law what would this equal?

Another name for this process is called **factoring**.

Example 4

Use the distributive law to factor the following:

a. $km + kn$

b. $5x + 10y$

There are a few other properties we need to mention:

The Additive Identity

For any number a , $0 + a = a + 0 = a$

The Multiplicative Identity

For any number a , $1 \times a = a \times 1 = a$

The Additive Inverse (aka the opposite of a number)

For any number a , the addition of its opposite will result in zero, ie: $a + (-a) = 0$.

The Multiplicative Inverse (aka the reciprocal)

For any number a , it's inverse can be written in the form $\frac{1}{a}$ and has the property: $a \times \frac{1}{a} = 1$

It is interesting to note that there exist number systems (not discussed in this class) where these properties do **not** exist!

Lesson: Like Terms and Combining Like Terms

In algebra, we often want to simplify expressions if possible to make our lives easier. This is where the discussion of like terms comes into play.

Like Terms

Like terms are terms that share the exact same variables with the exact same exponents. Only like terms may be added.

Do like terms work in real life? I.e., can you come up with an example of like terms that could still be added in the real world?

Building off of your example, what wouldn't be like terms and could NOT be added in that scenario?

Example 1

In the following list below, classify which terms are like terms.

Discuss the two examples with the class to ensure they understand how to identify like terms. Students may color code, underline, or draw shapes around anything in the lists to indicate they are like terms.

a. $2x, 7y, 6x^2, 8y, x, -2x^2$

b. $3x^2y^2, 7x^2y, 2xy^2, 8x^2y, x^2y^2, -3x^2, 5xy^2$

The idea of like terms only comes up when we are talking about addition or subtraction. We do not worry about this when we want to multiply two things together. You may only add or subtract terms that are like terms.

Example 2

Combine like terms. The next page is also left blank if you need more space.

a. $3x + 2y - 5x + 9y$

b. $4x^2 + 2x + 6y^2 + 2x^2 + 8x$

c. $\frac{1}{2}x + \frac{2}{3}y - \frac{3}{2}x - 2\frac{1}{3}y$

d. $3x^2y^3 + 7x^2y^2 - 2x^2y^3 + x^2y^2$

e. $\frac{1}{4}x + \frac{2}{5}y - \frac{1}{2} + \frac{1}{5}y$

Lesson: Translating English into Math

As we move through this course, one major goal is to learn how to recognize and interpret certain situations as math problems. There are particular words in English that denote certain operations in mathematics, and we will classify those here.

<u>Addition</u>	<u>Subtraction</u>	<u>Multiplication</u>	<u>Division</u>

Example 1

Translate the following sentences into an algebraic expression. There is nothing to solve for you, you are merely setting up expressions.

- | | |
|--------------------------------------|---|
| a. 3 more than Lucy's height | d. Five less than three times a number. |
| b. Two less than a number | e. Two-thirds of the cost. |
| c. Three more than twice his salary. | f. Three more than twice his and her salary combined. |

Rational Exponents

Video 1: How to work with rules like $x^{\frac{1}{n}}$ or $x^{\frac{m}{n}}$ or $x^{-\frac{m}{n}}$

In this video, we will learn how to evaluate expressions that look like this:

$$8^{\frac{1}{3}} \quad \text{or} \quad 25^{\frac{3}{2}}$$

Define $x^{\frac{1}{n}}$ and show how to use this rule to evaluate $25^{\frac{1}{2}}$

Examples Evaluate the following:

a. $125^{\frac{1}{3}}$

b. $32^{\frac{1}{5}}$

c. $9^{\frac{1}{2}}$

d. $(-9)^{\frac{1}{2}}$

e. $-9^{\frac{1}{2}}$

f. $\left(\frac{27}{125}\right)^{\frac{1}{3}}$

Define $x^{\frac{m}{n}}$ and show how to use this rule to evaluate $25^{\frac{3}{2}}$

Examples Evaluate the following:

a. $27^{\frac{2}{3}}$

b. $4^{\frac{3}{2}}$

c. $16^{\frac{5}{4}}$

d. $-8^{\frac{2}{3}}$

e. $(-8)^{\frac{2}{3}}$

f. $(-16)^{\frac{3}{4}}$

g. $-81^{\frac{3}{4}}$

h. $\left(\frac{125}{27}\right)^{\frac{2}{3}}$

Define $x^{-\frac{m}{n}}$ and show how to use this rule to evaluate $25^{-\frac{3}{2}}$

Examples Evaluate the following:

a. $25^{-\frac{1}{2}}$

b. $27^{-\frac{1}{3}}$

c. $16^{-\frac{3}{4}}$

d. $4^{-\frac{5}{2}}$

e. $\left(\frac{25}{4}\right)^{-\frac{3}{2}}$

f. $(-25)^{-\frac{3}{2}}$

g. $(-27)^{-\frac{2}{3}}$

h. $-16^{-\frac{5}{4}}$

Lesson: Using Exponent Rules with Rational Exponents

First, let's review the main exponent rules:

Next, let's do a quick review on operations with fractions. Complete the following operations:

a. $\frac{2}{3} \times \frac{5}{6}$

b. $\frac{2}{5} \times 10$

c. $\frac{2}{3} + \frac{2}{5}$

Now we will combine all of these ideas into the following examples!

Examples Simplify the following. Your final answer should not include any negative exponents.

a. $x^{\frac{2}{3}} * x^{\frac{1}{5}}$

b. $\left(4x^{\frac{2}{3}}y^6\right)^{\frac{3}{2}}$

c. $\frac{x^{\frac{2}{3}}y^{-\frac{1}{5}}}{x^{\frac{1}{6}}y^{\frac{2}{5}}}$

d. $\left(\frac{125x^4y^{-\frac{1}{3}}}{x^{-\frac{1}{2}}y^{\frac{2}{5}}} \right)^{-\frac{2}{3}}$

e. $\left(\frac{4x^6y^{-\frac{1}{3}}}{x^{-3}y^{\frac{1}{2}}} \right)^{-\frac{3}{2}}$

Lesson: Simplifying Radicals with Rational Exponents

Convert the following radicals to rational exponents, simplify, and then rewrite in radical form.

a. $\sqrt{7^2}$

b. $\sqrt{9^2}$

c. $\sqrt[3]{x^{15}}$

d. $\sqrt[4]{x^2}$

Rational Exponents

Video 3: Convert a radical expression to exponential form

Examples Convert the following expressions to exponential form, and then simplify if needed.

a. $\sqrt{7}$

b. $\sqrt[3]{8}$

c. $\sqrt{5^2}$

d. $\sqrt[3]{7^6}$

e. $\sqrt[4]{x^{16}}$

f. $\sqrt[6]{x^2}$

Lesson: Basic Exponent Rules – Product, Power, and Quotient Rules

When working with rules of exponents, it is very important to be able to identify the base of each exponent. To begin, we will practice this.

Example 1

State the base for each exponent in the problems below.

Expression	Exponent	Base for this Exponent
5^3		
-5^4		
$-5x^6$		
$3y^7$		

Product Rule

First we want to remember what the actual definition of an exponent means.

We want to find: $2^3 * 2^2$. We will calculate this a few ways to understand the rules.

First we will work towards finding the exact answer. Evaluate each exponent below as you normally would with the **order of operations**, and then multiply the numbers together:

Now we will re-examine this result solely from the viewpoint of exponents. Write out each exponent as a multiplication problem, and notice the results you get from this exercise.

What do you notice about the exponents?

Product Rule (formal mathematical definition)

Let x be a real number, and let a and b be integers. Then:

Example 2

Use the product rule to evaluate the following.

a. $x^2 * x^4$

c. $x^3 + x^7$

b. $3x^3 * -1x^4 * 2x$

d. $\frac{2}{3}x^4y^2z^5 * 6x^3y^2 * \frac{1}{2}y^3z^4$

Power Rule

Using the order of operations, first rewrite the expression inside the parentheses without exponents:

$(2^3)^2 =$

Now rewrite this entire result just utilizing what we know about the definition of exponents:

What is the relationship between the exponents in this case?

The Power Rule (formal mathematical definition)

Let x be a real number, and let a and b be integers. Then:

Example 3

Use the power rule to evaluate the following:

a. $(x^3)^5$

c. $(3x^4y^5)^3$

b. $(x^2y^7z)^4$

d. $\left(\frac{2}{3}x^5y^2\right)^3$

Is there a difference between $(x + y)^n$ and $(xy)^n$? Explain.

The Quotient Rule

Using the order of operations, first rewrite the expression inside the parentheses without exponents:

$$\frac{2^5}{2^3}$$

First we will evaluate and find the final answer by using the order of operations below.

Notice the final answer of this question. How can we rewrite this number in its exponent form?

What relationship do you see between the exponents in the original question and the final answer?

The Quotient Rule (formal mathematical definition)

Let x be a real number, and let a and b be integers. Then:

Example 4

Use the quotient rule to simplify the following:

a. $\frac{x^7}{x^3}$

c. $\frac{15x^4y^3z^{10}}{2x^2y^2z^5}$

b. $\frac{x^5y^8z^{10}}{xyz}$

d. $\frac{12x^7y^3}{8x^3z^4}$

Examples: Exponent Rules, Basic Examples

Simplify the following using only the product, power, and quotient rules. Your final answer should not contain negative exponents.

a. $x^2y^7 * 5x^3y^5z$

f. $(-3x^5)^2$

b. $(4x^7y^3)^2$

g. $(-3x^5)^3$

c. $\frac{15x^7y^3}{10x^2y}$

h. $(2x^4y^2z^6)^2 * 3x^2z^3$

d. $\frac{2}{5}x^2y^5 * \frac{1}{3}x^7yz^3$

i. $\frac{(x^3y^7)^2}{x^4y}$

e. $\left(\frac{5}{3}x^3y^2z^3\right)^3$

j. $\frac{(x^4y^2)^3}{(x^2y)^2}$

Examples: Exponent Rules, With Numbers Only

Simplify the following using only the product, power, and quotient rules. Your final answer should be a number.

a. $3^2 * 3$

d. $\left(\frac{2}{3}\right)^3 * 3^2$

b. $(-4)^2 * 2$

e. $\frac{6^2}{2^3}$

c. $\frac{2^7}{2^4}$

f. $\left(\frac{1}{2}\right)^4 * 2^2$

Examples: Exponent Rules, Trickier and Harder Examples

Simplify the following using only the product, power, and quotient rules. Your final answer should not include negative exponents.

a. $(3x^2y^4)^2 * (x^4y^5)^3$

d. $\left(\frac{3}{5}x^5y^2z^7\right)^2 * \frac{xy^3}{z^5}$

b. $\frac{(2x^5y^2)^3}{(x^2yz^5)^2}$

e. $\frac{(3x^5y^2z^4)^3}{(6x^2yz^3)^2}$

c. $\left(\frac{2}{3}x^5y^7z^3\right)^2 * 9x^3y^3$

f. $\frac{(4x^3y^2z^5)^2}{(2xz^2)^4}$

Lesson: Negative Exponents

Simplify: $\frac{25}{125}$

If we wrote these numbers as exponents, this calculation would look like this:

$$\frac{5^2}{5^3} = \frac{1}{5}$$

Consider what would happen if we used the quotient rule on this problem:

$$\frac{5^2}{5^3} = 5^{2-3} = 5^{-1} = \frac{1}{5}$$

This problem shows us that exponents can also be negative.

Three Rules for Negative Exponents

Let x and y be a real numbers, and let n be an integer. Then:

$$x^{-n} = \frac{1}{x^n}$$

$$\frac{1}{x^{-n}} = x^n$$

$$\left(\frac{x}{y}\right)^{-n} = \left(\frac{y}{x}\right)^n = \frac{y^n}{x^n}$$

In general your final answer cannot have a negative exponent, so it is essential that you understand what to do with it.

Example 1

Rewrite the following expressions without negative exponents.

a. 4^{-2}

c. $\left(\frac{2}{5}\right)^{-2}$

b. $\frac{1}{3^{-3}}$

d. $\left(\frac{1}{3}\right)^{-2}$

One of the biggest challenges with negative exponents is identifying how to apply them to the proper base. As a quick review, remember how we identified the bases in the last section:

Example 2

Identify the base of the following exponents:

Expression	First Base	First Exponent	Second Base	Second Exponent
3^5				
$-2x^5$				
$5x^3y^7$				

This is extremely important to remember when working with negative exponents. In general, our final answer can't have negative exponents so you need to be used to reformatting your answers properly if negative exponents come up.

Example 3

Simplify the following expressions, and write your final answer without negative exponents.

a. $\frac{a^3b^{-2}}{c^{-4}d^6}$

b. $\frac{3a^{-3}b^4c^{-5}}{d^2e^{-2}f^5}$

c. $\frac{a^5b^3c^4}{4d^{-2}e^{-4}f^{-2}}$

d. $\frac{a^{-3}b^{-2}c^{-6}}{d^4e^7f}$

With this in mind, we should now be able to work on combining all of the rules of exponents we've learned.

Example 4

Simplify the following. Your final answer cannot contain negative exponents.

a. $(3x^{-2}y^4)^3$

c. $(-2x^3y^{-2}z)^{-3}$

b. $\left(\frac{2x^{-4}y^6}{x^5y^{-3}}\right)^4$

d. $\left(\frac{2x^3y^{-4}}{3x^{-2}y^{-5}}\right)^{-3}$

Examples: Negative Exponents, Basic Examples

Your final answer should not contain negative exponents.

a. $\left(\frac{4}{5}\right)^{-2}$

e. $\frac{14x^{-3}y^5z^3}{21x^{-2}y^{-3}z^9}$

b. $3x^{-6}y^4z^{-2}$

f. $(3x^{-4}y^{-2}z^4)^2$

c. $\frac{4x^{-4}y^7}{z^{-5}}$

g. $\left(\frac{5x^2y^{-4}}{z^2}\right)^5$

d. $\frac{6x^3y^{-5}}{x^7y^2}$

h. $(x^{-3}y^2z^{-4})^{-3}$

Examples: Negative Exponents, Tricky Examples

Your final answer should not contain negative exponents.

a. $(3x^{-4}y^5z^{-2})^{-3}$

d. $\left(\frac{3x^3y^{-3}z^4}{2x^{-2}y^4z^{-2}}\right)^{-3}$

b. $(-5x^2y^{-3}z^4)^{-2}$

e. $\frac{(2x^{-4}y^5)^4}{x^{-3}y^7}$

c. $\left(\frac{4x^{-5}y^2}{x^{-3}y^{-5}}\right)^{-2}$

f. $\frac{(3x^{-3}y^2z^{-3})^{-3}}{(2x^4y^{-5}z^3)^{-4}}$

Examples: Negative Exponents versus Negative Numbers

Examples Simplify:

a. 6^{-2}

b. -6^2

c. $(-6)^2$

d. $(-6)^{-2}$

Examples Simplify:

a. $5x^{-2}$

b. $(5x)^{-2}$

c. $-5x^2$

d. $(-5x)^{-2}$

Examples Simplify

a. -7^2

b. $(-7)^{-2}$

c. 7^{-2}

d. -7^{-2}

Examples Simplify

a. $3x^{-2}$

b. $(-3x)^{-2}$

c. $(-3x)^{-3}$

Lesson: Scientific Notation

Scientific Notation

A number in scientific notation is of the form: $A \times 10^n$ where $1 \leq |A| < 10$ and n is an integer

Before we can write numbers in this form, it is best if we can identify examples that are or are not in this format.

Example 1

Identify if the following examples are in scientific notation. If not, circle and explain why.

- | | | |
|--------------------------------------|----------------------------------|-------------------------------------|
| a. 5.34×10^7 | Yes, it's in scientific notation | No, it's not in scientific notation |
| b. $-2.7832 \times 10^{\frac{1}{2}}$ | Yes, it's in scientific notation | No, it's not in scientific notation |
| c. -4.5382×10^{-8} | Yes, it's in scientific notation | No, it's not in scientific notation |
| d. 0.8×10^{23} | Yes, it's in scientific notation | No, it's not in scientific notation |

We will now examine how to rewrite the following example in scientific notation.

Example 1

Rewrite the following in scientific notation.

Example 2

Rewrite the following examples in their expanded form.

Example 3

Use the rules of exponents to multiply the following results together.

a. $(2 \times 10^6) \times (3 \times 10^8)$

b. $(2 \times 10^{17}) \times (2.1 \times 10^{23})$

c. $(8 \times 10^7) \times (2 \times 10^{-3})$

Example 4

Use the rules of exponents to divide the following results.

a. $\frac{8 \times 10^{12}}{4 \times 10^3}$

b. $\frac{6 \times 10^7}{2 \times 10^{19}}$

c. $\frac{3 \times 10^{-12}}{6 \times 10^{-5}}$

Example 5

Add the following:

a. $(5 \times 10^{78}) + (2 \times 10^{78})$

b. $(8 \times 10^6) + (2 \times 10^6)$

c. $(3 \times 10^4) + (2 \times 10^3)$

The Basics of Roots

Video 1: Square Roots

What are the square roots of 9? Write down what it means to be a square root.

Examples Find the square roots of the following numbers.

a.

b.

What does the symbol $\sqrt{\quad}$ refer to?

How can we refer to the negative square root?

Draw a square root and label its parts:

Examples Evaluate the following:

a. $\sqrt{25}$

b. $\sqrt{81}$

c. $\sqrt{36}$

d. $\sqrt{\frac{1}{36}}$

e. $\sqrt{\frac{16}{49}}$

f. $\sqrt{\frac{9}{25}}$

g. $\sqrt{-36}$

h. $-\sqrt{64}$

Finally, how would you estimate a root like $\sqrt{5}$?

Example Estimate $\sqrt{14}$

The Basics of Roots

Video 2: Higher Roots

How would you evaluate the following? Explain the rationale for how to evaluate these:

$$\sqrt[3]{8}$$

$$\sqrt[4]{81}$$

$$\sqrt[5]{32}$$

The most general symbol for a root is $\sqrt[n]{a}$. Explain what this symbol means and once again, label it.

Now we can create a list of the most likely roots that we will encounter. You will very likely want to create another list of this on a page you can easily access.

Squares

Cubes

Other

Examples Evaluate the following. Be careful when you evaluate.

a. $\sqrt[3]{125}$

d. $-\sqrt[3]{8}$

b. $\sqrt[3]{-125}$

e. $-\sqrt[3]{-8}$

c. $\sqrt[4]{16}$

f. $\sqrt[5]{-32}$

d. $\sqrt[4]{-16}$

g. $\sqrt[6]{-64}$

Now let's generalize. For all roots of the form $\sqrt[n]{a}$, what can we say if....

1. The index (n) is an **odd** number?
2. The index (n) is an **even** number and a is a **negative** number?
3. The index (n) is an **even** number and a is a **positive** number?

The Basics of Roots

Video 3: The “Ultimate” Test and A Few Other Notes

To begin, we will first see how well we can put everything we’ve learned so far about roots together.

Examples Evaluate the following, if possible.

a. $\sqrt[3]{-8}$

b. $\sqrt{-16}$

c. $-\sqrt[4]{16}$

d. $\sqrt[3]{-125}$

e. $-\sqrt{100}$

f. $\sqrt[4]{-81}$

g. $-\sqrt{25}$

h. $-\sqrt[3]{-8}$

i. $-\sqrt[4]{-16}$

j. $-\sqrt[5]{-32}$

And now that you have completed this ... it’s time for my “ultimate” test ... drum roll please:

The “Ultimate” Test for Roots

Evaluate. Note: at this moment in time, we do not know anything about imaginary numbers.

a. $\sqrt{1}$

h. $-\sqrt[3]{1}$

b. $\sqrt[3]{1}$

i. $-\sqrt{-1}$

c. $\sqrt[5]{1}$

j. $-\sqrt[7]{-1}$

d. $\sqrt[4]{-1}$

k. $-\sqrt[3]{-1}$

e. $\sqrt[3]{-1}$

l. $\sqrt[8]{-1}$

f. $-\sqrt{1}$

m. $-\sqrt[10]{-1}$

g. $-\sqrt[4]{-1}$

n. $\sqrt{-1}$

Are you having fun yet???

We have one last thing to discuss: the general rule for evaluating $\sqrt[n]{a^n}$

If n is even

If n is odd

Examples Evaluate the following:

a. $\sqrt[3]{y^3}$

c. $\sqrt[7]{(2y - 4)^7}$

b. $\sqrt[4]{(x - 4)^4}$

d. $\sqrt[6]{(3x + 2)^6}$

Simplifying Radicals

The Basics

We will look at how to simplify radicals like this:

$$\sqrt{50} = 5\sqrt{2} \quad \text{or} \quad \sqrt[3]{250} = 5\sqrt[3]{2}$$

You might want to have the list of squares / cubes / quartics ready for this exercise. If you don't, pause the video and write them all down on a separate sheet of paper so that you can have it handy for this entire lesson.

First we look at **the multiplication property of radicals**:

Examples Multiply the following:

a. $\sqrt[3]{2} * \sqrt[3]{3}$

b. $\sqrt{7} * \sqrt{5}$

c. $\sqrt[3]{3} * \sqrt[3]{9}$

d. $\sqrt[3]{6} * \sqrt[3]{9}$

How are examples c and d related?

Being able to multiply radicals together is powerful, because it also allows us to break apart radicals and simplify them. Using the example $\sqrt{12}$, write down how to simplify:

Examples Simplify the following:

a. $\sqrt{50}$

b. $\sqrt{20}$

c. $\sqrt[3]{54}$

d. $\sqrt[4]{32}$

e. $\sqrt[3]{24}$

At this point, we should probably define when a radical is actually simplified.

A simplified radical will

Examples Explain why the following radicals are not simplified:

a. $\sqrt{24}$

b. $\frac{5}{\sqrt{7}}$

c. $\sqrt[4]{\frac{3}{5}}$

d. $\sqrt[3]{8x^8y^2}$

Examples Simplify the following:

a. $\sqrt{24}$

b. $\sqrt[3]{32}$

c. $\sqrt[4]{64}$

d. $\sqrt[5]{64}$

e. $\sqrt{250}$

f. $\sqrt[3]{250}$

Lesson: Simplifying Radicals with Variables

We are going to rely on a skill we've already learned to help with simplifying radicals.

Examples Show you can simplify the following utilizing rational exponents.

a. $\sqrt{x^6}$

b. $\sqrt[3]{x^{15}}$

c. $\sqrt{x^6 y^8}$

d. $\sqrt[4]{x^{12} y^{20}}$

How can we simplify $\sqrt{x^5}$? Watch the 2 explanations in the video and summarize the explanation that makes more sense to you.

Example Simplify the following.

a. $\sqrt[3]{x^{13}}$

b. $\sqrt[4]{x^{22}}$

c. $\sqrt{12x^6y^7}$

d. $\sqrt[3]{250x^8y^{16}}$

e. $\sqrt[4]{\frac{32x^7}{y^8}}$

Lesson: Multiplying Radicals

Write a one-sentence explanation of how to multiply $\sqrt[3]{3} * \sqrt[3]{4}$

Examples Multiply the following. Simplify if possible.

a. $\sqrt[3]{6} * \sqrt[3]{4}$

b. $\sqrt[4]{4x^3} * \sqrt[4]{12x^2}$

c. $\sqrt{5x} * \sqrt{10xy}$

Simplifying Radicals

Multiplying and Dividing Radicals with Different Indices

We will look at how to perform operations like:

$$\sqrt{x} * \sqrt[3]{x} \quad \text{or} \quad \frac{\sqrt[4]{x}}{\sqrt[3]{x}}$$

Before we begin, you should be comfortable with doing problems like this:

$$8^{\frac{1}{3}}$$

$$25^{\frac{3}{2}}$$

Write down the approach for performing this operation: $\sqrt{x} * \sqrt[3]{x}$

Examples Perform the following operations:

a. $\sqrt[3]{x} * \sqrt[4]{x}$

b. $\frac{\sqrt[5]{x^2}}{\sqrt[3]{x}}$

Examples Perform the following operations:

a. $\sqrt[4]{x^3} * \sqrt[5]{x^3}$

b. $\frac{\sqrt{x}}{\sqrt[3]{x^2}}$

c. $\sqrt[3]{x^2} * \sqrt{x}$

Adding and Subtracting Radicals

In this video we will look at how to perform operations like:

$$\sqrt{3} + \sqrt[3]{5} - 5\sqrt{3} + 7\sqrt[3]{3} \quad \text{or} \quad \sqrt{24x^4y^3} - 3x^2y\sqrt{6y}$$

Before you begin, you should already be able to simplify radicals like this:

a. $\sqrt{24x^6y^9}$

b. $\sqrt[4]{16x^7y^{11}}$

What are like terms?

Add the like terms in this problem: $3x + 7y - 9x + 7y^2 - 2x - 2y$

What are like radicals?

Add the like radicals in this problem: $5\sqrt{3} - 2\sqrt[3]{3} + 8\sqrt[3]{3} - 2\sqrt{5} + 8\sqrt{3}$

Examples Perform the following operations:

a. $4\sqrt{7} - 5\sqrt[3]{3} + 2\sqrt{3} - 3\sqrt[3]{7} - 9\sqrt{3} + \sqrt[3]{3}$

b. $2x\sqrt{7} - 3\sqrt{7} + 8x\sqrt{7} - 2\sqrt{7}$

c. $3x\sqrt[3]{5} + 2x\sqrt[3]{4} - 3\sqrt[3]{4} + 5x\sqrt[3]{4} - 7x\sqrt[3]{5} + 5\sqrt[3]{4}$

It is possible that you might have to simplify a radical before you can add it. Note how this works with the example:

$$\sqrt{24} + 2\sqrt{54}$$

Examples Perform the following operations:

a. $5\sqrt{8} - \sqrt{18}$

b. $4\sqrt[3]{54} - 7\sqrt[3]{16}$

Examples Perform the following operations:

a. $\sqrt{8x^2y^4} - 9xy^2\sqrt{2}$

b. $4x^3\sqrt[3]{16x^6y^7} + 9x^3y^3\sqrt[3]{250y^4}$

c. $5x^2y^3\sqrt[4]{32x^7z^{10}} - 2x^3z^2\sqrt[4]{2x^3y^{12}z^2}$

d. $3x^3\sqrt[3]{24x^5y^7} + 3xy^3\sqrt[3]{3x^2y} + y^2\sqrt[3]{375x^8y}$

Lesson: Multiplying Radicals Expressions

First, let's review how to multiply radicals in general. Show how to multiply:

a. $\sqrt[3]{4x^2y} * \sqrt[3]{6x^2y}$

b. $\sqrt{6xy} * \sqrt{10y}$

Now we will apply that logic to larger rational expressions.

Write a one-sentence explanation of how to multiply: $\sqrt{2}(\sqrt{3} + \sqrt{10})$, then multiply.

Example Multiply $2\sqrt[3]{3}(\sqrt[3]{9} + 5\sqrt[3]{6})$

Example Multiply $3x\sqrt{5}(\sqrt{10x} - 5x\sqrt{2})$

Write out an explanation of how to multiply: $(\sqrt{3} - \sqrt{2})(\sqrt{6} + \sqrt{3})$

Example Multiply: $(3\sqrt{5} - \sqrt{2})(\sqrt{10} + 4\sqrt{2})$

Example Multiply: $(3\sqrt[3]{4} - \sqrt[3]{6})(5\sqrt[3]{2} + \sqrt[3]{6})$

How do you multiply $(\sqrt{2} + \sqrt{3})^2$

Examples Multiply: $(\sqrt[3]{9} + \sqrt[3]{6})^2$

Lesson: Rationalizing a Denominator with One Square Root

What does it mean to rationalize the denominator?

How would you rationalize $\frac{2}{\sqrt{3}}$? Explain *why* we can do this.

Can every fraction with a radical in the denominator be rationalized like this? Circle: Yes No

What types of problems can be rationalized this way?

Examples Rationalize the following.

a. $\frac{20}{\sqrt{8}}$

b. $\sqrt{\frac{40}{60}}$

c. $\sqrt{\frac{81x^7}{2y}}$

d. $\frac{\sqrt{22}}{\sqrt{x^9}}$

Dividing Radicals/Rationalizing the Denominator

Part 2: Higher Roots

In this video we will look at how to simplify expressions like:

$$\frac{5}{\sqrt[3]{3}} \quad \text{or} \quad \frac{\sqrt[4]{2}}{\sqrt[4]{9}}$$

Before you begin, you should already be able to simplify problems like this:

a. $\frac{5}{\sqrt{3}}$

b. $\frac{\sqrt{6}}{\sqrt{18}}$

First, let's observe why higher roots are a different situation. Write down what happens if we use the same methods as above to rationalize the following: $\frac{5}{\sqrt[3]{3}}$

We have to pivot our thinking to "count powers". Note how this works with this examples: $\sqrt[3]{3} * \underline{\hspace{1cm}} = 3$

Examples Determine what to fill in the blank to create a true statement:

a. $\sqrt[3]{25} * ? = 5$

e. $\sqrt[3]{7} * ? = 7$

b. $\sqrt[4]{2} * ? = 2$

f. $\sqrt[4]{25} * ? = 5$

c. $\sqrt[4]{27} * ? = 3$

g. $\sqrt[5]{3} * ? = 3$

d. $\sqrt[5]{4} * ? = 2$

h. $\sqrt[3]{36} * ? = 6$

Now you should have a sense of what number you want this to equal. This is a very strange exercise, but using the pattern above, determine what to multiply by to get to the desired root:

a. $\sqrt[3]{4} * ? =$

c. $\sqrt[5]{9} * ? =$

b. $\sqrt[4]{5} * ? =$

d. $\sqrt[4]{36} * ? =$

Now we should have a decent sense of how to rationalize and what the answer should equal. Explain how to rationalize the following:

a. $\frac{5}{\sqrt[3]{2}}$

b. $\frac{7}{\sqrt[4]{5}}$

Examples Rationalize the following denominators:

a. $\frac{6}{\sqrt[3]{5}}$

b. $\frac{2}{\sqrt[4]{9}}$

c. $\frac{2}{\sqrt[5]{125}}$

d. $\frac{7}{\sqrt[4]{8}}$

Examples Rationalize the following denominators:

a. $\frac{\sqrt[3]{5}}{\sqrt[3]{4}}$

d. $\frac{6\sqrt[3]{5}}{\sqrt[3]{3}}$

b. $\frac{2\sqrt[3]{3}}{\sqrt[3]{4}}$

e. $\frac{3}{\sqrt[5]{x^3}}$

c. $\frac{5\sqrt[4]{2}}{\sqrt[4]{27}}$

f. $\frac{\sqrt[5]{25}}{\sqrt[5]{16}}$

Lesson: Rationalizing a Denominator with Two Square Roots

How do you find the conjugate of $4 + \sqrt{2}$? Write one sentence to explain what you need to do.

Examples What is the conjugate of the following?

a. $3 - \sqrt{5}$

b. $-4 - \sqrt{2}$

Show how to rationalize: $\frac{5}{\sqrt{2}-\sqrt{3}}$

Examples Rationalize the following.

a. $\frac{\sqrt{2}}{\sqrt{3}-\sqrt{6}}$

b. $\frac{\sqrt{2}-\sqrt{3}}{\sqrt{3}+\sqrt{5}}$

c. $\frac{\sqrt{x}-\sqrt{y}}{\sqrt{x}+\sqrt{y}}$

Lesson: Factoring by GCF and Trinomials (the more memorable types of factoring)

How do you factor $3x + 3y$?

What type of factoring is this called?

Examples Show how to use the GCF to factor the following:

a. $4x + 12x^2 - 8x^3$

b. $5x^2y + 10x^7y^3 - 15x^5y^2$

Examples Show how to use the GCF to factor the following:

a. $4x^2 + 12xy - 3y^2$

b. $15x^3y^4 - 25x^6y^2 + 20x^7y^8$

What is a trinomial?

Show how to multiply: $(x + 2)(x + 7)$

Show how to factor $x^2 + 3x + 2$ using the *trial and error* method. Write down some reasoning explaining how this method works.

Example Factor the following using trial and error.

a. $x^2 + 5x + 6$

b. $x^2 - 7x + 12$

c. $x^2 - 25$

Show how we can factor $2x^2 + 5x + 3$ using *trial and error*.

Examples Factor the following.

a. $5x^2 + 7x + 2$

b. $3x^2 + 13x - 10$

Lesson: Factoring by Grouping and the Sum and Difference of Cubes

Show how to use grouping to factor: $ax + ay + bx + by$

What are some key hints that you probably need to use the grouping technique?

Write out the steps of how to use grouping for this problem:

Example Show how to finish the factoring for the following:

a. $5(x - 2y) + z^2(x - 2y)$

b. $6x^2(3y - z) - (3y - z)$

Examples Use grouping to factor the following:

a. $4x^3 - 12xy + x^2y - 3y^2$

b. $xy + 4x - y - 4$

For the next example we will use $12x^2 - 15y + 18x - 10xy$

What is the issue when we try to use grouping?

How can we resolve this issue and still use grouping?

Write the sum and difference of cubes formulas below:

Examples Factor the following:

a. $x^3 + 8$

b. $y^3 - 1$

c. $8x^3 + 27y^3$

Lesson: A Few More Tricks for Factoring

List all the types of factoring that you might have to use:

It is always possible that you will have to combine these methods. There are also a few other tricks you can try, which we will discuss in this lesson.

Using **Grouping** with Trinomials

Write out *instructions* on how to use grouping for $4x^2 + 16x + 15$

Below are a few challenging examples of factoring. Write out any notes of how to work these out that might be helpful to you. Remember, you are expected to use these notes as a resource to help you finish your homework – take the time to write down reminders that will be meaningful and helpful to you!

a. $-x^2 - 15x - 36$

What trick can you use here? Show how to use it.

b. $4y^3 - 28y^2 + 48y$

What trick can you use here? Show how to use it.

c. $-6x^2 + 10x - 3$

d. $x^2 + 7xy + 12y^2$

How should you deal with having the multiple variables?

Finally, let's work with a few trickier examples of the difference of squares. Follow the pattern:

a. $x^2 - 16$

b. $x^2 - 25$

c. $x^2 - \frac{16}{25}$

d. $16 - x^2$

e. $25x^2 - 16y^2$

f. $\frac{16}{25}x^2 - \frac{4}{9}y^2$

Lesson: A Review of Exponent Rules for Algebra Students

Power Rule: $x^m * x^n =$

Product Rule: $(x^m)^n =$

Quotient Rule: $\frac{x^m}{x^n} =$

Negative Exponents

$$x^{-n} =$$

$$\frac{1}{x^{-n}} =$$

$$\left(\frac{x}{y}\right)^{-n} =$$

Simplify the following using exponent rules. Your final answer should not contain negative exponents.

a. $3x^7y^{-4} * 5x^{-2}y^9$

e. $\frac{8x^{-3}y^4z^{-3}}{10x^5y^{-2}z^{-7}}$

b. $\left(\frac{2x^4y^5}{3z^5}\right)^3$

f. $\left(\frac{3x^{-3}y^{-2}}{6x^2y^4}\right)^{-2}$

c. $\frac{10x^4y^7}{2x^2y^3z^{-5}}$

g. $\left(\frac{2x^{-7}z^3}{3x^{-2}y^{-3}z^5}\right)^{-3}$

d. $\frac{4x^3y^{-2}}{6x^{-4}y^{-5}}$

h. $\left(\frac{5x^{-2}y^{-3}z^3}{x^3y^4z^{-7}}\right)^{-2}$

Lesson: An Intro to Polynomials

We can extend our knowledge of exponents to another algebraic topic – polynomials

Before we define what a polynomial is, it is usually easier to see a few examples of what is or is not a polynomial.

The following ARE polynomials:

- $x^2 - 3x + 8$
- $x^7 - 5x^4 + 2x^3 + x$
- 9
- $7x + \pi$
- $x^3 + \frac{2}{3}$

The following ARE NOT polynomials:

- $\sqrt{x} - 4$
- $4x^{-4} - 7x^{-3}$
- $\frac{4}{x} - \frac{2}{x+3}$

Now we can more formally define a polynomial.

Polynomial

An expression with real numbers, variables, and whole number exponents.

Descending Order

Ordering a polynomial from the largest exponent to the smallest exponent, and any freestanding number without a variable is last

Example 1

Put the polynomials in descending order.

a. $x^3 + 4x^5 - 2x^7 + 3x^2 + 5x^8$

b. $x^2 - x^3 + x^4 - x^5 - x^6 + 3$

Term

Any part in a polynomial that is separated by a “+” or “-” sign is considered to be its own term

Variables

The letters in a polynomial that represent numbers

Coefficients

The numbers attached to the variables

Constant Term

The freestanding number within a polynomial that is not attached to any variables

Example 1

Examine the polynomials, then determine the variables, coefficients, and constant terms

Polynomial	Terms	Variable(s)	Coefficient(s)	Constant
$2x + 3$				
$3x^2 - x + 5$				
5				
$2x^2y + 3y - x$				

Degree of a Term

The degree of a term is the sum of the exponents of that term.

Example 2

Identify the degree of each term

Term	Degree	Term	Degree
x^4		x^4y^3	
$3y^6$		x^7y^9	
x		$x^4y^2z^5$	
6		xy^4	

Degree of a Polynomial

The degree of a polynomial is the same as the degree of the term of highest degree within that polynomial

Leading Coefficient

The leading coefficient is the coefficient attached to the term of highest degree in a polynomial

Trinomial

A polynomial with 3 terms

Binomial

A polynomial with 2 terms

Monomial

A polynomial with 1 term

Example 3

Identify the degree of each polynomial and the type of polynomial it is.

Polynomial	Degree of Each Term, Respectively	Deg of Polynom	Leading Coefficient	Type of Polynomial
$x^5 + 4x^3 - 2x$				
$-5x^2 + 18x$				
$6x + 2x^4 - 3x^2$				
$x^5 + 2x^7 - 4$				
8				

Evaluating Polynomials

Evaluating polynomials is not unlike evaluating other expressions we've seen in this course.

Example 4

Evaluate the polynomials for $x = -2$.

a. $x^2 + 3x + 1$

b. $x^3 - x^2 - 3x + 4$

c. $\frac{1}{4}x^3 - \frac{1}{2}x^2 + 5x$

Lesson: How to Add and Subtract Polynomials

To add or subtract polynomials, combine like terms.

Examples

Perform the following operations.

a. $(2x + 4) + (5x^2 - 3x + 1)$

b. $(5x^2 - 3x + 1) - (x^3 - 2x^2 - x + 1)$

c. $(3x^3 + 2x^2 - 5x) + (4x^3 + 5x^2 + 2x)$

d. $(2x^2 + 5x - 4) - (x^3 - 2x^2 + 3x - 5)$

Lesson: How to Multiply Polynomials

To begin, let's first focus on the process of distributing a monomial with a polynomial.

Examples Multiply the following:

a. $3x(4x^2 + 5x - 7)$

b. $x^2y^4(3x^2 + 2xy - 8y^4)$

c. $\frac{1}{2}x^3y^4(4xy^3 - 2x^3y + 8xy)$

Now write down the process for how to multiply $(x + 2)(x^2 + 3x)$

Examples Multiply the following:

a. $(x + 3)(x + 2)$

b. $(3x + 1)(x - 6)$

c. $(x^2 + 4)(x - 2y)$

d. $(x + 4)(x - 4)$

e. $(x + 3)(x^2 + 4x + 5)$

f. $(2x^2 + 3x + 1)(3x^2 - 4x + 2)$

How do you multiply $(x + 5)^2$?

Examples Multiply the following:

a. $(x + 2)^2$

b. $(2x^2 + 1)^2$

c. $(3x^2 + x + 5)$

How do you multiply 3 polynomials together, such as $(x + 2)(x + 1)(x - 3)$?

Examples Multiply the following together:

a. $(x - 3)(x + 4)(x - 1)$

b. $(2x + 1)(x - 2)(x - 1)$

Lesson: The Basics of Dividing Polynomials

In this section, we will discuss how to divide polynomials that are in these forms:

$$\frac{x^2+3x}{x} \quad \text{or} \quad (4x^7y^3 + 8x^4y^2 - 12x^5y^5) \div 2x^2y$$

To divide in these cases, you can just use rules of exponents. Sometimes you will want to separate the problem into separate fractions.

Examples

a. $\frac{x^2+3x}{x}$

b. $\frac{5x^4-10x^3+15x^2}{5x}$

c. $(27x^4y^2 + 18x^3y) \div 9x^2y$

d. $(4x^7y^3 + 8x^4y^2 - 12x^5y^5) \div 2x^2y$

Lesson: Long Division for Polynomials

In this section, we will discuss how to divide polynomials that are in these forms:

$$(x^2 + 5x + 6) \div (x + 3) \quad \text{or} \quad (2x^2 + 5x + 3) \div (2x + 3)$$

In these types of problems, we need to use long division. In another video, we talked about how to divide polynomial problems like these:

$$\frac{4x^3 + 2x^2}{2x}$$

But we did not need long division. What is different about these types of problems that require long division?

Show how to use long division to divide $(x^2 + 5x + 6) \div (x + 3)$

Examples Divide the following.

a. $(x^2 + 7x + 10) \div (x + 5)$

b. $(2x^2 + 5x + 3) \div (2x + 3)$

c. $\frac{x^2 - 11x + 30}{x - 5}$

Now we will look at what to do when there is a remainder.

Divide the following: $(x^2 + 3x + 1) \div (x + 2)$

Now show how to divide when there are “missing” terms, such as with $(x^4 - 81) \div (x^2 - 9)$

Examples Divide.

a. $(5x^2 - 10x^3 + 8 - 9x) \div (x + 2)$

b. $(x^3 + 1) \div (x + 1)$

c. $(5x^3 - 4x^2 + 1) \div (x + 2)$

A Quick(ish) Review of Factoring

We will quickly go over the 4 types of factoring in this video.

- Greatest Common Factor (GCF)
- Grouping
- Trinomials
- Sum/Difference of Cubes

Greatest Common Factor

Show how the GCF works with the example: $3x + 3y$

Examples Factor the following.

a. $15x^2y + 10xy^3$

b. $27xy^4 - 18x^2z + 81x^4y^3$

c. $7xy - 3z^4$

Grouping

Show how to use grouping with: $ax + ay + bx + by$

a. $3x + 6y - xy - 2y^2$

b. $x^4 - x^2 - 3x^3 - 3x$

Trinomials

What are some of the standard forms of trinomials presented in the video?

Examples Factor using trial and error.

a. $x^2 + 3x + 2$

b. $x^2 + 7x + 10$

c. $-x^2 + 2x + 15$

d. $x^2 - 4$

e. $x^2 + 5xy + 6y^2$

f. $2x^2 + 5x + 3$

Sum and Difference of Cubes

Write down the formulas for the sum and difference of cubes.

Examples Factor.

a. $x^3 + 27$

b. $8x^3 - y^3$

Lesson: The Greatest Common Factor

The Greatest Common Factor (GCF)

The Greatest Common Factor is the biggest common factor between a set of terms.

Examples

Find the greatest common factor in the following polynomials, then factor the polynomial.

a. Polynomial: $3x + 3y$

GCF:

Factor out the GCF:

b. Polynomial: $3xy + 6xz$

GCF:

Factor out the GCF:

c. Polynomial: $10x^2 - 25xy + 15xz$

GCF:

Factor out the GCF:

d. Polynomial: $3x^7y + 8x^3z^5 - 5y^4z^3$

GCF:

Factor out the GCF:

Examples

Factor the following polynomials.

a. $12xy^2 + 10x^2y^4 - 14xy^5$

b. $5xy - 10yz + 12xz$

c. $3x + 6x^2 - 12x^3$

d. $5x^4y^2z^5 + 10x^2y^4z^6 - 25x^5yz^3$

Lesson: Factoring by Grouping

Grouping can best be understood with the following example.

Factor $ax + ay + bx + by$ by grouping:

Examples

Finish the following examples where the grouping had been started.

a. $3(x - y) + 2z(x - y)$

b. $2a(3b + 2c) - 5(2c + 3b)$

c. $4x(y - z) - 3(z + y)$

d. $4z(4x - 7) - (4x - 7)$

Examples

Factor the following.

a. $5x + 10y + xz + 2yz$

b. $10x^2 + 15x^3y - 2 - 3xy$

c. $3xy + 4y - 27x - 36$

d. $3xy + 6x + 21y + 42$

e. $3x^2 - 48y + 8x - 18xy$

Lesson: The Basics of Factoring Trinomials of the Form $x^2 + bx + c$

Factoring trinomials will require us to multiply polynomials, so first we will review this skill.

Examples

Multiply.

a. $(x + 3)(x - 5)$

b. $(2x - y)(5x + 2y)$

Definition of a Trinomial

A trinomial is a polynomial with three terms.

How to Factor a Trinomial Using Trial and Error

Show how to factor $x^2 + 3x + 2$

Show how to factor $x^2 + 7x + 12$

Examples

Factor the following.

a. $x^2 + 6x + 5$

b. $x^2 + 12x + 11$

c. $x^2 - 8x + 15$

d. $x^2 - 13x + 30$

e. $x^2 + 10x - 24$

f. $x^2 - 5x - 14$

g. $2x^2 + 16x + 30$

h. $5x^4 + 55x^3 - 45x^2$

Lesson: The Basics of Factoring Trinomials of the Form $-x^2 + bx + c$

How can you factor $-x^2 - 5x - 6$?

What about $-x^2 - 15x - 36$?

Examples

Factor the following.

a. $-x^2 + 13x - 30$

b. $-x^2 + 16x - 55$

c. $-x^2 - 2x + 3$

Lesson: The Basics of Factoring Trinomials of the Form $x^2 + bxy + cy^2$

How do you factor $x^2 + 5x + 6$?

How do you factor $x^2 + 5xy + 6y^2$?

Examples Factor the following.

a. $x^2 + xy - 12y^2$

b. $x^2 - 2xy + y^2$

c. $3x^2 + 21xy + 18y^2$

Lesson: The Basics of Factoring Trinomials of the Form $ax^2 + bx + c$

In this lesson we will focus on trinomials that do not have a leading coefficient.

To begin, first let's get warmed up by factoring the following using trial and error:

a. $x^2 + 7x + 12$

b. $x^2 - 3xy + 10y^2$

How do you factor $2x^2 + 5x + 3$ using trial and error?

How do you factor $3x^2 + 16x + 5$ using trial and error?

Another way to factor $3x^2 + 16x + 5$ is by grouping. Show how to use this technique below:

Examples Factor the following:

a. $7x^2 - 11x + 4$

b. $3x^2 + 10x + 8$

c. $5x^2 - 11x - 36$

d. $3x^2 - 17xy - 6y^2$

e. $-10x^2 + 19x - 6$

Lesson: Factoring Special Trinomials

How do you factor $x^2 + 6x + 9$?

What's another way to write this factorization?

What do you call this type of trinomials?

Examples Factor the following:

a. $x^2 + 12x + 36$

b. $9x^2 - 12x + 16$

c. $x^2 - x + \frac{1}{4}$

d. $36x^2 - 60xy + 25y^2$

How do you factor $x^2 - 4$?

What is this type of factoring called?

Examples Factor the following:

a. $x^2 - 25$

b. $x^2 - 16$

c. $x^2 - 1$

d. $25x^2 - 36$

e. $16x^2 - 81y^2$

f. $16 - x^2$

g. $25y^2 - 64x^2$

h. $\frac{49}{64} - x^2$

Lesson: How to Factor The Sum and Difference of Cubes

Write down the formula for the sum of cubes

$$a^3 + b^3 =$$

Write down the formula for the difference of cubes

$$a^3 - b^3 =$$

Show how to factor $x^3 + 8$

Show how to factor $8x^3 - 27$

Examples Factor:

a. $x^3 - 1$

b. $x^3 - \frac{1}{8}$

c. $x^3 + 64$

Review: An Overview of All the Types of Factoring

We will review all of the different types of factoring by presenting an example of each type.

Greatest Common Factor

1. $4x^2 + 8xy$

2. $6x^3y^2 - 8x^2y^5 + 10x^5y$

Grouping

3. $3x + 6y - xz^2 - 2yz^2$

4. $12x^2 - 2y + 3x - 8xy$

Basic Trinomials of the Form $x^2 + bx + c$

5. $x^2 + 7x + 10$

6. $x^2 - 2x - 15$

Trickier or Special Forms of Trinomials of the Form $x^2 + bx + c$

7. $-x^2 - 3x + 28$

8. $6x^2 + 54x + 120$

9. $x^2 - 6xy + 8y^2$

10. $x^2 - 8x + 16$

11. $x^2 - 25$

12. $x^2 - \frac{9}{4}$

Trinomials of the Form $ax^2 + bx + c$

13. $2x^2 + 11x + 5$

14. $7x^2 - 15x + 8$

15. $-3x^2 - 15x^2 + 198x$

16. $16x^2 - 24xy + 9y^2$

More Practice with the Difference of Squares

17. $x^2 - 81$

18. $25x^2 - 4$

19. $\frac{1}{4}x^2 - 9y^2$

Sum and Difference of Cubes

20. $x^3 + 8$

21. $x^3 - 27$

22. $27x^3 - 1$

23. $8x^3 + 64y^3$

Rational Expressions

Video 1: The Basics

What are rational expressions?

What do rational expressions look like? Write down some examples:

Simplifying Rational Expressions

Take down some notes on this:

Examples together:

a. $\frac{25x^7y^9}{15x^3y^2z^7}$

Examples on your own:

b. $\frac{21x^7y^9}{14y^8z^3}$

c. $\frac{16x^4y^7z^3}{15x^3y^4z^8}$

Examples together:

a. $\frac{x^2+5x+6}{x^2-9}$

Examples on your own:

b. $\frac{x^2+7x+10}{x^2+10x+25}$

c. $\frac{x^2-x-6}{x^2-9x+14}$

d. $\frac{x^2+10x+16}{x^3+2x^2+x+2}$

Play the video and check your work.

Examples on your own:

e. $\frac{x^3-8}{x^2-9x+14}$

f. $\frac{10x^3+15x^7}{10x^3+40x^2+40x}$

Rational Expressions

Video 2: Other Things to Know

When it comes to fractions, where are you allowed to have a zero? In the numerator (top) or denominator (bottom)?

Therefore, what do we have to worry about with rational expressions?

Examples on your own:

Evaluate the following rational expressions for $x = 2$.

a. $\frac{x^2+6x+8}{x^2+8x+15}$

b. $\frac{x^2+9x+20}{x^2+3x-10}$

Example on your own: (evaluate for x=2)

c. $\frac{x^2-4}{x^2+9x+14}$

To summarize...

When a rational expression has a zero in the numerator, what does this mean?

When a rational expression has a zero in the denominator, what does this mean?

Define the term **domain**:

What are we concerned with for the domain of rational expressions?

A review of interval notation:

Examples:

Write the following in interval notation:

a.

b.

c.

Example together:

Determine i) where the expression equals zero ii) where the expression is undefined iii) the domain of the expression

a. $\frac{x+3}{x-5}$

b. $\frac{2x+1}{x^2+7x+10}$

Examples on your own:

Determine i) where the expression equals zero ii) where the expression is undefined iii) the domain of the expression

c. $\frac{x-3}{x-1}$

d. $\frac{x^2-81}{x^2+5x+6}$

Rational Expressions

Video 3: The Negative One Trick with Simplifying

In its current form, can you simplify: $\frac{x-3}{3-x}$? Why or why not?

Show how you can manipulate this expression to simplify it.

There is no formal name for this trick. Some might refer to it as a *multiplicative property of -1*, but I refer to as “that -1 trick.” I know, it’s not exactly formal sounding!

Let’s do one more example of this. Simplify $\frac{5-x}{3x-15}$

The first thing to get used to it how to rewrite expressions using this trick.

Examples

Rewrite the order of each expression by factoring out -1 first.

a. $4 - x$

b. $3x - 1$

c. $x - 6$

Now let's use this to simplify rational expressions.

Examples Simplify the following.

a. $\frac{6x-12}{2-x}$

Examples Simplify the following.

b. $\frac{4x-20}{30x-6x^2}$

c. $\frac{x^2-4}{2-x}$

d. $\frac{x-6}{36-x^2}$

e. $\frac{5-x}{x^2-7x+10}$

Examples Simplify the following.

f. $\frac{x^2-10x+25}{25-x^2}$

g. $\frac{27-x^3}{x^2-5x+6}$

h. $\frac{x^3-3x^2+5x-15}{6-3x}$

Examples Simplify the following.

i. $\frac{54-9x}{x^2-36}$

j. $\frac{8-x^3}{x^2-9x+14}$

k. $\frac{y^3-x^3}{x^2-y^2}$

Example/Question: Troubleshooting the -1 Trick

Explain the actual DETAIL behind WHY $\frac{4}{5-x} = -\frac{4}{x-5}$

Examples Write an equivalent fraction for the following:

a. $\frac{5}{x-8}$

b. $\frac{x-3}{x+2}$

Lesson: Multiplying Rational Expressions

How do you multiply the fractions $\frac{a}{b} \times \frac{c}{d}$? Include a one-sentence explanation.

Examples Multiply the following.

a. $\frac{4}{5} \times \frac{7}{9}$

b. $\frac{2x^3}{7y^2} \times \frac{5x^5}{9y^7}$

Show how to make cancellations before multiplying $\frac{12}{25} \times \frac{15}{16}$

We can apply this same logic to rational expressions.

Examples Multiply the following and simplify when possible.

a. $\frac{21x^7}{15y^3} \times \frac{5y^8}{42x^4}$

b. $\frac{2x^2+5x+3}{x^2-25} \times \frac{x^3+125}{4x^2+6x}$

Lesson: Dividing Rational Expressions

How do you divide $\frac{a}{b} \div \frac{c}{d}$? Write a one sentence explanation of how to do this too.

Example Divide the following: $\frac{5x}{9} \div \frac{2}{7x^4}$

When can you make cancellations in the process?

Examples Divide.

a. $\frac{12x^7}{45} \div \frac{42x^3}{15}$

b. $\frac{x^2+4x+4}{x+5} \div (x+2)^5$

c. $\frac{x^3+1}{6x^3-6x^2+6x} \div \frac{x^2-1}{8}$

Explain why $\frac{\frac{5x+5y}{24}}{\frac{15x^2+15xy}{12}}$ is both a fraction and a division problem.

Example Divide: $\frac{\frac{16x^2-25}{x^5}}{\frac{36x-45}{9x^7}}$

Lesson: Finding the Least Common Denominator for Addition and Subtraction – A Quick Review for Fractions (with only numbers)

We are going to work on adding and subtracting rational expressions with different denominators. Before we do that, we are going to remind ourselves how to find common denominators for fractions that only have numbers.

The simplest case: what is the least common denominator between the fractions $\frac{1}{2}$ and $\frac{1}{3}$?

Why is that the common denominator?

Now let's look at something slightly more complicated. What is the least common denominator between $\frac{1}{6}$ and $\frac{1}{10}$?

First, take a guess. It's totally fine if this guess is wrong, but just make a guess.

Why is this your guess? What did you do?

There is a specific procedure that you want to use when finding least common denominators in this class. Write down instructions that were given in the video, but write them out in a way that makes sense to you.

Repeat this procedure to find the least common denominator $\frac{1}{15}$ and $\frac{1}{10}$.

Repeat this procedure to find the least common denominator $\frac{1}{12}$ and $\frac{1}{8}$.

Now we will focus on how to write fractions with this desired denominator.

The simplest case: using the least common denominator you found for $\frac{1}{2}$ and $\frac{1}{3}$, show how to write 2 equivalent fractions with the desired denominator. Write out instructions on how you do this. Summarize what was discussed in the explanation in the video.

Using the least common denominator you found for $\frac{1}{6}$ and $\frac{1}{10}$, show how to write 2 equivalent fractions with the desired denominator. Write out instructions on how you do this. Summarize what was discussed in the explanation in the video.

Using the least common denominator you found for $\frac{1}{12}$ and $\frac{1}{8}$, show how to write 2 equivalent fractions with the desired denominator. Write out instructions on how you do this. Summarize what was discussed in the explanation in the video.

Lesson: Finding the Least Common Denominator for Rational Expressions

The simplest case: what is the least common denominator between the fractions $\frac{1}{2x}$ and $\frac{1}{3y^2}$?

Examples Find the least common denominator between the following fractions.

a. $\frac{1}{2x^2}, \frac{1}{3xy^2}$

b. $\frac{1}{2x^2y^5}, \frac{1}{3x^7y^2}$

c. $\frac{1}{2x^3y^5z^2}, \frac{1}{3x^5y^7z^8}$

Examples Find the least common denominator for the following.

a. $\frac{1}{14x^3y^2}, \frac{1}{21x^2y^7}$

b. $\frac{1}{24x^3y}, \frac{1}{12x^2y^5}$

What is the least common denominator between $\frac{1}{x+2}$ and $\frac{1}{x}$?

What is the least common denominator between $\frac{1}{x^2-4}$ and $\frac{1}{x^2+5x+6}$?

Examples Find the least common denominator between the following fractions.

a. $\frac{3}{5x-10}, \frac{1}{7x-14}$

b. $\frac{1}{x^2-12x+35}, \frac{1}{x^2+x-20}$

Examples Find the least common denominator between the following fractions.

a. $\frac{1}{x-3}, \frac{1}{3-x}$

b. $\frac{1}{x^2-4}, \frac{1}{2-x}$

c. $\frac{1}{x^2+9x+20}, \frac{1}{x^2-25}, \frac{1}{x^2-x-20}$

Lesson: Writing Equivalent Rational Expressions with the Same Denominator

Examples Find the least common denominator between the following rational expressions, and then write each rational expression with its equivalent and the LCD.

a. $\frac{2}{15x^2y^5}, \frac{4}{9x^3y^2}$

b. $\frac{5}{x+3}, \frac{4}{x}$

Examples Find the least common denominator between the following rational expressions, and then write each rational expression with its equivalent and the LCD.

a. $\frac{x}{x^2+9x+18}, \frac{2}{x^2+5x+6}$

b. $\frac{x}{5x^2-30x}, \frac{x-2}{2x^2-17x+30}, \frac{2}{2x^2-5x}$

Lesson: Adding and Subtracting Rational Expressions

Show how to add or subtract the following:

a. $\frac{5}{x+2} - \frac{3}{x+2}$

b. $\frac{5}{x-4} + \frac{7}{x}$

c. $\frac{5}{12xy^2} - \frac{7}{8x^3y}$

d. $\frac{x}{x+3} - \frac{2}{x^2-9}$

e. $\frac{x}{x^2-25} - \frac{5}{x^2-3x-40}$

Example/Question: When Are You Allowed to Cancel with Rational Expressions?

What is the wrong way to cancel $\frac{(x+1)-3(x-3)}{2(x-3)+4(x+1)}$ and explain why you can't cancel like that? How can you tell it's wrong?

What is the difference between $\frac{(x+1)-3(x-3)}{2(x-3)+4(x+1)}$ and $\frac{3(x+1)(x-3)}{2(x-3)(x+1)}$?

How do you simplify $\frac{(x+1)-3(x-3)}{2(x-3)+4(x+1)}$?

Lesson: Complex Fractions

First let's review how to simplify a complex fraction we've already discussed. Show how to simplify:

$$\frac{\frac{2x}{x+5}}{\frac{2}{x^2-25}}$$

Explain what makes the problem below different from the one we just reviewed above.

$$\frac{\frac{1}{2} - \frac{1}{3}}{\frac{5}{6} + \frac{1}{3}}$$

Outline the steps required to simplify. Write out the instructions as they were explained in the video.

$$\frac{\frac{1}{2} - \frac{1}{3}}{\frac{5}{6} + \frac{1}{3}}$$

Example Simplify the following

$$\frac{\frac{7}{9} - \frac{1}{3}}{1 + \frac{5}{9}}$$

Examples Simplify the following.

a.
$$\frac{\frac{5}{x} + \frac{2}{y}}{\frac{1}{y} - \frac{3}{x}}$$

b.
$$\frac{\frac{\frac{6}{x+3} - \frac{4}{x-1}}{2} + \frac{5}{x+3}}{x-1}$$

Example Simplify the following.

$$\frac{1 - \frac{4}{t+5}}{\frac{4}{t^2-25} + \frac{t}{t-5}}$$

Complex Fractions

Video 2: More Complicated Examples

Complex fractions look like this:

$$\frac{\frac{4x}{15y^2}}{\frac{6x^4}{25y^8}} \quad \text{or} \quad \frac{\frac{1}{3} - \frac{1}{6}}{\frac{3}{2} + \frac{5}{6}} \quad \text{or} \quad \frac{\frac{4}{x^2} - \frac{4}{y^2}}{\frac{1}{xy} + \frac{5}{y}}$$

First we will discuss the basics.

Examples

a. $\frac{\frac{\frac{1}{x^2} - \frac{1}{y}}{3 - \frac{4}{x}}}{y^2}$

b. $\frac{\frac{\frac{3}{x} - x}{\frac{2}{x} + x}}{x}$

Examples

a.
$$\frac{\frac{3}{x-1} - \frac{2}{x+3}}{\frac{5}{x+3} - \frac{2}{x-1}}$$

b.
$$\frac{\frac{5}{x-2} - \frac{3}{x+5}}{\frac{1}{x+5} + \frac{2}{x-2}}$$

Examples

a.
$$\frac{1 - \frac{1}{x+5}}{\frac{x}{x-5} + \frac{4}{x^2-25}}$$

b.
$$\frac{1 + \frac{12}{x^2-16}}{\frac{x}{x+4} - \frac{6}{x-4}}$$

c.
$$\frac{\frac{1}{x+2} + \frac{4}{x^2-4}}{\frac{3}{x-2} + 1}$$

Lesson: A Quick Review of Some Things About Graphing

Remind yourself how to graph the following points. Draw your own graph.

Point 1: $(-3, 4)$

Point 2: $(0, -2)$

Draw another rectangular coordinate system below and label the quadrants and origin.

Often when graphing, we use equations in two variables.

Example Find the ordered pair associated with the equation $3x + y^2 = 7$ when $x = 1$

Example Find at least 3 ordered pairs for the following equations.

a. $y = \sqrt{x + 2}$

b. $2x + 3y = 6$

Finding 3 points can be a strategy for graphing.

Example Sketch a graph for the previous 2 examples.

What is the x-intercept? And how do you find it?

What is the y-intercept? And how do you find it?

Examples Find the x- and y-intercepts for the following equations.

a. $y = 3x - 1$

b. $x = \sqrt{y - 1}$

c. $x^2 - 3x + 2 = y$

Lesson: The Distance Formula

Graph the points $(2, 4)$ and $(-4, -6)$. Explain how you can find the distance between these points by summarizing the explanation given in the video (make a decent effort to explain this – you will have an easier time remembering the formula if you understand WHY it works).

State the distance formula:

Example Find the distance between the following points: $(4, -1)$ and $(7, 3)$

There are 2 ways we can use the distance formula.

Example Show how to use the distance formula to determine whether the following points are the vertices of a right triangle.

a. $(0, -2)$, $(-4, 3)$ and $(5, 1)$

b. $(-2, -8)$, $(0, -4)$ and $(-4, -7)$

What does it mean for 3 points to be collinear?

Example Show how to use the distance formula to determine if the 3 points are collinear.

a. $(-1, 4), (-2, -1), (1, 14)$

b. $(-2, -4), (-6, 11), (0, -12)$

Lesson: The Midpoint Formula

What is the midpoint formula?

So what is the midpoint formula effectively doing?

Example Find the midpoint between the points $(-5, -6)$ and $(-3, 2)$

Example The midpoint of a line segment is $(5, 8)$ and the endpoint of that line segment is $(11, 10)$. What is the OTHER endpoint on this line segment?

Example The midpoint of a line segment is $(3, 1)$ and the endpoint of that line segment is $(6, 8)$. What is the OTHER endpoint on this line segment?

Example The midpoint of a line segment is $(2a, 5b)$ and the endpoint of that line segment is $(-1a, 9b)$. What is the OTHER endpoint on this line segment?

Lesson: Equations of Circles and How to Graph Circles

Draw the circle in the video and label the relevant parts of it.

Using your sketch above, explain how the distance formula can be used to help us determine the equation of a circle (you will need to summarize the explanation in your own words).

What is the **Center-Radius Form** of the equation of a circle?


Example Find the center radius form of a circle using the given information.

a. center $(-2, 3)$ and radius 4


b. center $(0, -1)$ and radius 2

Example Graph the following circles.

a. $(x - 1)^2 + (y + 3)^2 = 4$



b. $x^2 + y^2 = 9$



Expand $(x - 1)^2 + (y + 3)^2 = 4$ and show how this creates the general form.

What is the **General Form** of the equation of a circle?

Example Review how to use completing the square with the equation: $x^2 - 6x + 4 = 0$.

Example Show how to use completing the square to write each in the center-radius form, and then sketch a graph of the circle.

a. $x^2 - 4x + y^2 + 8y - 44 = 0$

b. $x^2 + y^2 + 2x - 6y + 14 = 0$

Examples: Working with the General Form Equation of a Circle

Example Show how to use completing the square to write in the center-radius form, and then sketch a graph of the circle if possible.

a. $4x^2 + 4y^2 + 4x + 16y - 19 = 0$

b. $x^2 + y^2 - 4x - 4y + 8 = 0$

c. $x^2 + y^2 + 2x - 4y + 18 = 0$

d. $9x^2 + 9y^2 + 12x - 18y - 23 = 0$

Examples: Graphing Circles Using Technology

You can use this page to take notes for how to solve the following 2 examples:

- a. Find all coordinates of points whose distance from $(-3, 2)$ and $(-2, 1)$ is $\sqrt{5}$.
- b. There's an earthquake! There are 2 receiving stations with the following coordinates:

$$A = (-1, 1), B = (-2, 1), C = (-3, -1)$$

The epicenter of the earthquake is 2 units from stations A and C, and $\sqrt{5}$ units from B. Where is the epicenter?

Lesson: Solving Linear Equations, A Review of the Basics

To begin, let's review some of what we've learned about working with equations and using graphs to solve them.

Linear Equation in One Variable

An equation that can be written in the form $ax+b=0$ where a and b are real numbers

Solution

A value that when substituted into the variable of an equation makes the equation true

Example 1

Determine if $x = -1$ is a solution to the following equations.

a. $5 - 4x = 9$

b. $2x + 3 = 1 - 3x$

We will now work with some basic ideas of solving linear equations. There are two key concepts we must understand to do this.

The Addition Principle

Let a , b , and c be real numbers. If $a=b$ then:

Example 1

Use the addition principle to solve the following:

Choose at least one from each category. Label the examples you choose as a, b, c, etc to keep the class organized.

a. $x - 4 = 2$

b. $x + 3 = 3$

c. $x - \frac{1}{3} = \frac{1}{2}$

The Multiplication Principle

Let a, b, c be real numbers. If $a=b$ then:

Example 2

Use the multiplication principle to solve the following:

a. $3x = 15$

b. $5x = 0$

c. $-\frac{2}{5}x = 10$

Example 3

Solve the following examples:

a. $5 - 4x = 33$

b. $3x + 2 = 2 + 4x - 8$

c. $3(4 - 2x) = 3x + 6$

Lesson: Solving Linear Equations, No Solution and Infinite Solutions

There are 2 special outcomes we have to watch out for:

An Example Where There Is No Solution

$$3(2x - 1) = 2(2x + 1) - 2(3 - x)$$

An Example Where There Are Infinite Solutions

$$4x + 5 = 3(2x + 1) - 2(x - 1)$$

A Summary of the 3 Possible Solution Types for Linear Equations

Examples: Solving Linear Equations, Basic Examples with No Fractions or Decimals

a. $3x - 5 = 7$

e. $4x + 7 + 6x = 7(1 - 2x)$

b. $6 - 2x = 6$

f. $3(5 - 2x) - 6(3 - 2x) = 5(1 + 2x)$

c. $4x + 7 = 2x - 9$

g. $4(2x - 1) - 2(6 - 3x) = 2(3x - 4)$

d. $3 - 4x - 7 + 2x = 4 - 6x - 8 + 4x$

h. $3(2x - 5) - 4x + 1 = 4(x - 3) - 2(1 - 3x)$

Lesson: Solving Linear Equations with Fractions

To begin, we will look at 3 examples to demonstrate how to clear fractions without parentheses. Take notes as shown in the video.

a. $\frac{1}{2}x + \frac{1}{5} = \frac{6}{5}$

b. $\frac{1}{2}x + \frac{1}{5} - \frac{1}{10}x = 1 + \frac{3}{10}x - \frac{2}{5}$

c. $\frac{1}{2}x + 1 + \frac{1}{4}x = \frac{3}{4} + \frac{2}{3} + \frac{1}{3}x$

Now we will look at how to clear fractions in equations with parentheses.

a. $\frac{3}{4}(3 + 2x) - \frac{1}{2}(5 - 3x) = \frac{5}{4}$

b. $\frac{1}{2}(x + 2) + \frac{1}{3}(2x - 5) = \frac{1}{3}(4 + 2x)$

Examples: Solving Linear Equations with Fractions (No Parentheses)

a. $\frac{1}{5}x - \frac{1}{3} = \frac{2}{3}$

b. $\frac{1}{3}x - 1 + \frac{1}{5}x + \frac{2}{5} = \frac{2}{5}x + \frac{1}{5}$

c. $\frac{1}{4}x - \frac{1}{10} + \frac{1}{5}x - \frac{4}{5} = \frac{3}{20}x + \frac{1}{10} + \frac{3}{10}$

d. $\frac{2}{3}x + \frac{1}{3} - \frac{1}{2}x = \frac{5}{6}x - \frac{1}{6} - \frac{2}{3}x + \frac{1}{2}$

Examples: Solving Linear Equations with Fractions (With Parentheses)

a. $\frac{1}{2}(3x - 2) - \frac{2}{5}(2x - 3) = \frac{1}{5}(x + 1)$

b. $\frac{3}{4}(4x + 5) + \frac{1}{2}(2 - 3x) = \frac{3}{4}(2x + 1)$

c. $\frac{1}{2}(2x + 3) - \frac{1}{3}(2x - 1) = \frac{1}{4}(x + 7)$

d. $\frac{3}{4}(2x - 1) + \frac{5}{8}(2 - 4x) = \frac{1}{4}(2 - 4x)$

Lesson: Solving Linear Equations with Decimals

To begin, we will look at 3 examples to demonstrate how to clear decimals without parentheses. Take notes as shown in the video.

a. $0.03x + 0.04 = 0.19$

b. $0.05x + 0.1 - 0.03x = 0.08$

Now we will look at 3 examples to demonstrate how to clear decimals with parentheses. Take notes as shown in the video.

a. $0.4(x + 2) - 0.3(2x + 1) = 0.09$

b. $0.1x + 0.03(4 - 2x) = 0.5(x + 3)$

Examples: Solving Linear Equations with Decimals (no parentheses)

a. $0.1x - 0.06 - 0.2x - 0.04 = 0.3$

b. $0.3x - 0.6 + x - 0.2 = 0.6x - 0.8 + 0.7x$

c. $1 + 0.2x - 0.7 + 2x = 0.5 - 0.4x + 1.8 + 0.6x$

Examples: Solving Linear Equations with Decimals (with parentheses)

a. $0.4(3x + 1) - 1.5(2x + 1) = 0.6(3 - 3x)$

b. $0.2(3x + 1) - 0.05(2x + 8) = 0.1(2x + 4)$

c. $0.09(2 - 4x) - 0.1(2x - 1) = 0.2(2 - x) - 0.06(2 + 6x)$

Lesson: Solving Rational Equations

How do you know that you need to solve a problem versus simplify it?

Explain how to solve the rational equation. Include some instructions in your notes.

$$\frac{2}{x-1} + \frac{1}{x+4} = \frac{4}{x+4}$$

What do we have to watch out for when solving rational equations?

Examples For the following, which solutions would NOT work for these rational equations?

a. $\frac{4}{x+1} - \frac{2}{x+3} = \frac{5}{x+7}$

b. $\frac{4}{x^2-9} + \frac{3}{x-3} = \frac{2}{x+3}$

Examples Solve.

a. $1 - \frac{8}{x} = \frac{10}{x}$

b. $\frac{5}{x-3} - \frac{2}{x^2-9} = \frac{5x+13}{x^2-9}$

c. $\frac{x^2}{2} = \frac{x^2-6x}{3}$

d. $\frac{4}{x+1} = \frac{10}{x^2-1} - \frac{5}{x-1}$

Solving Equations with Rational Expression

We will learn how to solve equations that look like this:

$$z = \frac{x+3y}{4w} \quad \text{or} \quad \frac{1}{a} - \frac{2}{b} = \frac{3}{c}$$

Examples

a. Solve for y: $z = \frac{3xy}{5w}$

b. Solve for x: $w = \frac{7+2y}{xz}$

c. Solve for z: $y = \frac{4xw}{z-w}$

Examples

a. Solve for x: $\frac{3}{x} + \frac{1}{y} = \frac{2}{z}$

b. Solve for b: $\frac{1}{a} - \frac{1}{b} = \frac{3}{c}$

c. Solve for z: $\frac{5}{x} - \frac{3}{y} = \frac{7}{z}$

Lesson: Three Ways to Use the LCD with Rational Expressions

We have spent quite a bit of learning how to find the LCD between rational expressions and then using it in very different ways. We are going to compare the 3 techniques for how we used the LCD in this video.

First, simplify completely:

$$\frac{9}{x+5} - \frac{4}{x+1}$$

Before we begin, should x have a solution? Why or why not?

Now show your work below.

What type of problem is this? How did we leverage the LCD in this case?

Simplify completely:

$$\frac{\frac{9}{x+5} - \frac{4}{x+1}}{\frac{10}{x^2+6x+5} - \frac{3}{x+1}}$$

Before we begin, should x have a solution? Why or why not?

Now show your work below.

What type of problem is this? How did we leverage the LCD in this case?

Solve:

$$\frac{9}{x+5} - \frac{4}{x+1} = \frac{10}{x^2 + 6x + 5}$$

Before we begin, should x have a solution? Why or why not?

Now show your work below.

What type of problem is this? How did we leverage the LCD in this case?

Example/Question: When Can You Cross Multiply

Explain what a ratio is:

When can you cross-multiply?

Solve: $\frac{8}{x+2} - \frac{12}{x-4} = \frac{2}{x+2}$. Show that you CANNOT cross-multiply in this situation.

Now show why cross-multiplying works with $-\frac{x}{5} = \frac{3}{x+8}$

Lesson: Applications and Modeling with Linear Equations

We will revolve a few different types of applications for linear equations.

Example Two sides of a triangle have the same length. The third side measures 8 inches less than twice the common length. The perimeter of the triangle is 20 inches. What are the lengths of the three sides?

Example Dan ran the 100-m dash with a time of 9.92 seconds. If this pace could be maintained for an entire 26-mile marathon, what would his time be? (Note: 1 m = 3.281 ft, 1 mi = 5280 ft)

Example Sam earned \$25,000 from royalties on his music. He set aside 20% of this for a down payment on a new home. The balance will be used for a new car (eventually). He invested a portion of the money in a CD account (bank certificate of deposit) that earns 2.25%, and the remainder in another CD account that earns 1.75%. If the total interest earned after 1 year is \$405, how much money was invested in each account?

Lesson: Applications of Quadratic Equations

Example Find the problem of 2 pairs of 2 consecutive integers whose product is 156.

Example You want to buy a rug for a room that is 18ft x 26ft. You want to have a uniform strip of floor around the rug, and you can afford 384 square feet of carpeting. What dimensions should the rug have?

Example The area of a square is 32 more than its perimeter. Find the length of a side.

Lesson: The Basics of Applications and Word Problems

As a reminder, here is a review of some of the key words that imply certain operations in math:

<u>Words That Imply Addition</u>	<u>Words That Imply Subtraction</u>	<u>Word That Imply Multiplication</u>	<u>Words That Imply Division</u>
Add	Subtract	Multiplied by	Divided by
Increased by	Difference	Times	Quotient of
Altogether	Less than	Of	Out of
Sum	Fewer	Product of	Ratio
Longer than	Decrease	Twice	
Plus	Minus		
Above	Take away		
More than	Shorter than		

Example Juan is memorizing a list of names for a history class quiz. So far, he has memorized 8 of the names, which is one-fourth of the list. How many names are on this list?

Example Corey wants to rent a truck for the day. It costs \$30 for the base fee, plus \$0.23 for each mile driven. When he returned the truck, he owed \$41.19. How many miles did he drive?

Example A café serves both tea and coffee. On Sunday, the café sold 95 teas and coffees combined. The café sold 9 more coffees than teas. How many teas were sold on Sunday?

Example Latoya bought a new sweater and a pair of jeans for \$64. The jeans cost \$8 less than the sweater. How much was the sweater? How much were the jeans?

Example There are red, orange, and green balls in a ball pit. There are 12 more orange balls than red balls, and there are 9 more green balls than red balls. How many are there of each color?

Lesson: Consecutive Integer Word Problems

How do you represent consecutive integers with variables?

Example The sum of 2 consecutive integers is 73. Find the integers.

Example The sum of 3 consecutive integers is -33. Find the integers.

How do you represent even consecutive integers with variables?

How do you represent odd consecutive integers with variables?

What is noteworthy about how you represent even or odd consecutive integers?

Example The sum of 2 consecutive even integers is 110. Find the integers.

Example The sum of 3 consecutive odd integers is -87. Find the integers.

Example Find 3 consecutive odd integers such that their sum is seven more than four times the largest integer.

Lesson: Solving a Linear Equation with Multiple Variables

Examples

a. Solve for x: $x + y = 7$

b. Solve for y: $4y = z$

c. Solve for a: $3a - 6b = 2c$

d. Solve for b: $4a + 8b = 16c$

e. Solve for k: $\frac{1}{3}k = m$

f. Solve for a: $3(a - 2b) = 6c$

g. Solve for b: $\frac{2}{3}(3a + 2b) = c$

h. Solve for a: $-\frac{2}{5}(2a - 4b) = 4$

Lesson: Ratios and Proportions

What is a ratio?

Example The unit price of an item is the ratio of the price of an item to the amount of the item. Find the unit price if 5oz of peanuts costs \$1.

Example Express the ratio of 12 boys to 14 girls.

What is a proportion?

Examples Solve the following proportions:

a. $\frac{x}{5} = \frac{6}{10}$

b. $\frac{x+2}{12} = \frac{x-4}{8}$

Examples: Proportions

a. $\frac{3}{x} = \frac{9}{21}$

b. $\frac{3x+10}{15} = \frac{2}{5}$

c. $\frac{2x-1}{6} = \frac{4x}{10}$

Lesson: Proportion Word Problems

It 3 cookies cost \$1.36, how much will 7 cookies cost?

At a motorcycle race, the ratio of parents to children was 5 to 3. If 500 children attended the motorcycle race, how many parents were at the race?

One euro is equivalent to \$1.18 US dollars. How many euro would you get for \$25?

Lesson: Distance, Rate, and Time Problems

Two cars leave Glen Ellyn, IL, with one driving north and the other driving south. The northbound car is traveling 8mph faster than the southbound car, and after 2 hours they are 420 miles apart. Find the speed of each car.

Marshall and Lilly live together. Marshall leaves the house at 2pm, and then Lilly leaves the house at 2:30pm. If Marshall is driving 30 mph and Lilly is driving 35 mph in the same direction, how long will it be until their cars meet?

A Lamborghini and a Ferrari pass one another at noon on a highway in California. The cars are going opposite directions. If the Lamborghini is traveling 75 mph, and the Ferrari is traveling 80 mph, how long will it take for the cars to be 310 miles away from one another?

Lesson: Money Word Problems

Sarah has five more nickels than quarters. When combined, the coins are worth \$4.75. How many nickels and quarters does she have?

Maria has \$75 worth of \$1 and \$5 bills. In total, she has 27 bills. How many \$5 bills does she have?

Tickets at the movie theater course \$15 for an adult and \$10 for a child. In the last hour, sales were \$550. How many adult tickets versus child tickets were sold?

Lesson: Applications of Quadratic Equations

The area of a rectangle is 15ft squared, and the length of the rectangle is 7ft less than twice the width. Find the dimensions of the rectangle.

An 12ft ladder leans against the side of a house. The bottom of the ladder is 6ft from the side of the house. How high is the top of the ladder from the ground? If necessary, round your answer to the nearest tenth.

A kite flying in the air has a 14ft line attached to it. Its line is pulled taut and casts a 9ft shadow. Find the height of the kite. If necessary, round your answer to the nearest tenth.

The longer leg of a right triangle is 2in longer than the shorter leg. The hypotenuse is 4in longer than the shorter leg. Find the side lengths of the triangle.

Solving Formulas with Squares and Square Roots

In this lesson, we will look at how to solve for formulas that look like this:

$$y = 3x^2 \quad \text{or} \quad M = \sqrt{\frac{AB}{C}} \quad \text{or} \quad kx^2 + cx - n = 0$$

Solving Formulas with Squares

Solve the following for the indicated letters:

a. for x: $y = \frac{4x^2}{z}$

b. for k: $b = \frac{3}{k^2}$

Solving Formulas with Square Roots

Solve the following for the indicated letters:

a. for b: $a = \sqrt{\frac{b}{c}}$

b. for z: $x = \sqrt{\frac{3y}{z}}$

Solving Formulas in other forms

Solve the following for the indicated letters:

a. for x : $kx^2 + cx - d = 0$

b. for b : $ab^2 - 5b + m = 0$

c. for k : $\frac{1}{2}k^2 - 5nk + m = 0$

Solving Applications of Quadratic Equations

An object is thrown upwards off a cliff that is 48 feet above the ground. The height of the object t seconds after the object is released is given by:

$$h = -16t^2 + 32t + 48$$

How long does it take the object to hit the ground?

How long does it take the object to reach a height of 20 feet?

A 12 ft by 16 ft pond is going to be surrounded by a border of grass of uniform width all the way around. If the area of the enclosure must be 480 feet squared, what is the width of the grass border?

How to Evaluate Square Roots that Have Imaginary Number Answers and Dealing with Negatives in Square Roots

In this video, we will look at how to evaluate things like this:

$$\sqrt{-4} \quad \text{or} \quad \sqrt{-12} \quad \text{or} \quad \sqrt{-3} * \sqrt{-6}$$

What is an imaginary number?

What is a complex number?

How do you evaluate something like $\sqrt{-9}$?

Examples Compute the following:

a. $\sqrt{-25}$

b. $\sqrt{-81}$

c. $\sqrt{-12}$

d. $\sqrt{-7}$

e. $\sqrt{-\frac{25}{16}}$

f. $\sqrt[3]{-8}$

g. $\sqrt[4]{-16}$

h. $\sqrt{-32}$

How do you evaluate problems like $\sqrt{-3} * \sqrt{-2}$?

Examples Evaluate the following:

a. $\sqrt{-6} * \sqrt{-5}$

d. $\frac{\sqrt{-6}}{\sqrt{-2}}$

b. $\sqrt{-6} * \sqrt{-2}$

e. $\frac{\sqrt{-50}}{\sqrt{-2}}$

c. $\sqrt{-10} * \sqrt{-15}$

f. $\frac{\sqrt{-7}}{\sqrt{-2}}$

Operations with Imaginary Numbers

In this video, we will look at how to evaluate things like this:

$$2i(3i - 5) \quad \text{or} \quad (4i + 1) - (5i + 3) \quad \text{or} \quad \frac{4}{3+2i}$$

Adding and Subtracting with Imaginary Numbers:

Use the example $(2i + 3) + (5i - 6)$ to show how to add or subtract with imaginary numbers.

Examples Simplify:

a. $(3i - 2) - (4i - 9)$

b. $(5i + 3) + (2i - 1)$

What can we say about i^2 ?

With that in mind, explain what is unique about multiplying: $2i(3i + 1)$

Examples Simplify:

a. $4(2 - 3i)$

b. $6i(3i + 2)$

c. $5i(2 - 8i)$

d. $(5i + 1)(2i - 3)$

e. $(4i - 3)(6i + 7)$

Examples Simplify:

a. $(5i + 2)^2$

b. $(3i - 5)^2$

Why do you need to rationalize $\frac{3}{2i+1}$? Explain how to do this.

Examples Rationalize the following:

a. $\frac{4}{2i+6}$

b. $\frac{7}{i+3}$

Examples Rationalize the following:

a. $\frac{3i+1}{5i-2}$

b. $\frac{7i+3}{2i+1}$

c. $\frac{i-3}{5i+3}$

Evaluating Higher Powers of Imaginary Numbers

In this video, we will look at how to evaluate things like this:

$$i^{21} \quad \text{or} \quad i^{54}$$

First we need to investigate the powers of i and the patterns with it. Write this down below:

Now that we can see a pattern, explain how to evaluate the following:

$$i^{15}$$

$$i^{22}$$

Examples Evaluate the following:

a. i^{36}

b. i^{43}

c. i^{52}

d. i^{33}

e. i^{25}

Lesson: A Review of Imaginary Numbers

Please note: Knowing imaginary numbers is one of the “prerequisite” topics for this course. Ie – you should already know this (though I also realize you probably need a review!). If you have *never* seen imaginary numbers before, then I strongly suggest that you watch my series on imaginary numbers (because, yes, I have one). So if you felt like this went too fast and you want more details, please take responsibility for your education and go back to review this!

How do we represent an imaginary number?

What is i^2 ?

What is a complex number?

When are 2 complex numbers equal?

What does $\sqrt{-a}$ mean?

Examples Write using the definition of $\sqrt{-a}$

a. $\sqrt{-12}$

b. $\sqrt{-25}$

Examples Complete the following operations.

a. $\sqrt{-5} * \sqrt{-3}$

b. $\frac{\sqrt{-12}}{\sqrt{-18}}$

c. $\frac{\sqrt{-50}}{\sqrt{9}}$

To add or subtract complex numbers, we can add/subtract the real parts and the imaginary parts similar to working with like terms.

Examples Find the sum or difference.

a. $(2 - 6i) + (8 + 2i)$

b. $(5 - 7i) - (-4 + 2i)$

When multiplying complex numbers, recall that $i * i = i^2 = -1$!

Examples Multiply.

a. $(2 + 4i)(3 - 5i)$

b. $(4 - 5i)(3 - 2i)$

c. $(3 + 2i)(3 - 2i)$

Recall, the conjugate is found by using the opposite side for the second term!

Examples What is the conjugate of the following?

a. $4 + 2i$

b. $-3 - 4i$

c. i

To find the quotient of 2 complex numbers, we multiply by the conjugate of the denominator.

Examples Find each quotient. Write your answer in *standard form*, ie $a + bi$ form

a. $\frac{3-4i}{2+i}$

b. $\frac{5}{i}$

How do the powers of i cycle through?

Examples Show how to simplify the following powers of i . Write your work in great detail so that you remember how to do this!

a. i^{17}

c. i^{-14}

b. i^{28}

d. i^{-5}

Examples The following quadratic equations have imaginary solutions. Show how to solve them.

a. $x^2 = -25$ using the **square root property**

b. $x^2 + 75 = 0$ using the **square root property**

c. $2x^2 - 3x = -6$ using the **quadratic formula**

d. $x^4 - 16$ using **factoring** and also the **square root property**

Lesson: A Quick Review of How to Solve Quadratic Equations

What is the form of a quadratic equation?

How is this different from a quadratic function?

What are the 4 ways you can solve a quadratic equation?

Which technique should you use when solve a quadratic equation?

We will review these 4 techniques on the next page!

What type of answers are possible when you solve a quadratic equation?

Example Show how to use each of the 4 different techniques for solving quadratic equations by using each one at most once for the following examples.

a. $5x^2 + 6x + 1 = 0$

b. $(x + 4)^2 = 12$

c. $x^2 - 4x + 8 = 0$

d. $x^2 - 5x + 1 = 0$

Optional extra help: If you need a more thorough (and more complete) review of the methods we discussed today, watch these:

- A review of ALL the different types of factoring (just factoring, nothing about solving quadratic equations): <https://youtu.be/nrl7dJRxe5I> and <https://youtu.be/yGY5eqZkJcw>
- How to solve a quadratic equations by factoring: <https://youtu.be/5uof6OEyod0>
- How to solve quadratic equations by the square root property: <https://youtu.be/a1-smI3Xx5w>
- How to solve quadratic equations by completing the square: <https://youtu.be/6jXpMTozmhl>
- How to solve quadratic equations by the quadratic formula: <https://youtu.be/ryp2rdkhks8>
- A review that compares all the techniques on the same problem: <https://youtu.be/2ny9HbQ8doY>

A Quick Refresher of the 4 Ways to Solve Quadratics

For all of these examples, we will solve $x^2 + 6x + 5 = 0$.

Solving by Factoring

Solving by Completing the Square

Solving by Using the Quadratic Formula

The quadratic formula is: $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

One final way we can solve quadratics is by using the square root property.

Use the square root property to solve $(x + 3)^2 = 12$

Choosing the Best Method to Solve a Quadratic

There are 4 ways we can solve quadratics:

- | | |
|----|----|
| 1. | 3. |
| 2. | 4. |

Using each of these methods only once, determine the best way to solve these quadratics:

- a. $x^2 + 6x = 2$
- b. $x^2 + 3 = 2x$
- c. $(2x + 1)^2 = 49$
- d. $3x^2 + 2 = 3x$

Examples Using each of these methods only once, determine the best way to solve these quadratics:

a. $(3x - 2)^2 = -24$

b. $2x^2 + 5x - 1 = 0$

c. $x^2 - 12 = x$

d. $x^2 - 2x + 5 = 0$

Lesson: A Quick Review of The Discriminant

What is the discriminant?

Fill in the table below:

If the discriminant is ...	How many solutions are there?	What type of solution?
Positive and a perfect square		
Positive (not perfect square)		
Zero		
Negative		

Example Use the discriminant to determine how many solutions there are to: $x^2 - 4 = 0$.

Example Use the discriminant to determine how many solutions there are to: $x^2 + 6x + 9 = 0$.

Lesson: Solving for Quadratic Variables

Example Solve for t: $s = \frac{1}{2}gt^2$

Examples Solve for x: $kx^2 - tx = m$

Solving Quadratic Equations with Factoring

In this video, we will look at how to solve equations by factoring, which looks like this:

$$x^2 + 5x + 6 = 0 \rightarrow (x + 2)(x + 3) = 0$$

Quadratic equations are equations of the form:

Where a , b , and c are real numbers and $a \neq 0$.

How would you solve the quadratic equation $x^2 + 7x + 6 = 0$ by factoring?

Examples Solve the following by factoring:

a. $x^2 - x - 12 = 0$

b. $x^2 + 9 = -6x$

c. $x^2 = 25$

Examples Solve the following by factoring:

a. $x^2 + 7x = 0$

b. $\frac{1}{10}x^2 + \frac{1}{2}x + \frac{3}{5} = 0$

c. $2x^3 + 28x^2 + 90x = 0$

Solving Quadratic Equations using the Square Root Property

In this video, we will look at how to solve equations by using the square root property, which looks like this:

$$(x + 2)^2 = 9 \rightarrow x + 2 = \pm 3$$

Quadratic equations are equations of the form $ax^2 + bx + c = 0$ where a, b, and c are real numbers and $a \neq 0$.

What are the solutions to $x^2 = 25$?

In general, how do you solve equations of the form $Q^2 = k$, where Q is an expression and k is a constant? In other words, what is the **square root property**?

Examples Solve the following equations using the square root property:

a. $(x + 3)^2 = 25$

b. $(x + 3)^2 = 12$

Examples Solve the following equations using the square root property:

a. $(5x - 2)^2 + 5 = 29$

b. $(6x + 7)^2 + 4 = -41$

Solving Quadratic Equations by Completing the Square

In this video, we will look at how to solve equations by factoring, which looks like this:

$$x^2 + 6x = 7 \quad b = 6, \quad \frac{1}{2}b = 3, \quad \left(\frac{1}{2}b\right)^2 = 9$$

Quadratic equations are equations of the form $ax^2 + bx + c = 0$ where a , b , and c are real numbers and $a \neq 0$.

Before we begin, you should be familiar with the square root property. Solve $(2x + 1)^2 = 49$ using the square root property.

First let's talk about how to complete the square. Show how to complete the square on $x^2 + 6x$ below:

Examples Complete the square on the following:

a. $x^2 + 8x$

b. $x^2 - 10x$

c. $x^2 + 3x$

d. $x^2 - \frac{1}{2}x$

e. $x^2 + \frac{2}{3}x$

Show how to solve $2x^2 + 12x - 16 = 0$ by completing the square:

Examples Solve the following by completing the square:

a. $x^2 + 4x - 6 = 0$

b. $x^2 + 5 = 2x$

Examples Solve the following by completing the square:

a. $3x^2 + 15x = -6$

b. $\frac{1}{2}x^2 + 3x - 2 = 0$

Examples: The Distance Formula

Find the distance between the following sets of points.

a. $(6, 2)$ and $(2, 11)$

b. $(3, -1)$ and $(-4, -2)$

Solving Quadratic Equations by the Quadratic Formula

Complete the square on: $ax^2 + bx + c = 0$

State the quadratic formula:

Examples Use the quadratic formula to solve the following:

a. $x^2 + 3x + 2 = 0$

b. $2x^2 + 3x = 5$

c. $\frac{1}{2}x^2 + \frac{1}{3}x + \frac{5}{6} = 0$

Lesson: The Discriminant

Restate the Quadratic Formula, and then label the discriminant.

Value of the Discriminant	Interpretation

Examples Find the values of the discriminant for the following equations. Then solve the equations (using the most efficient method) to determine if this makes sense.

a. $x^2 - 4x + 4 = 0$

b. $x^2 + 6x + 5 = 0$

c. $3x^2 + 2x + 5 = 0$

Solving Equations that End Up Having Quadratics

In this lesson, we will review some techniques from previous lessons.

Solve $\frac{2}{x} - \frac{1}{x+1} = \frac{2}{3}$

Solve $\sqrt{2x + 8} = x + 4$

Examples Solve the following:

a. $\frac{4}{x-2} + \frac{x}{x-3} = \frac{2}{x-3}$

b. $\sqrt{3x+5} = x-3$

Solving Equations that Are Similar to Quadratics

In this lesson, we will look at how to solve equations that look like this:

$$x^{\frac{2}{3}} - x^{\frac{1}{3}} - 12 = 0 \quad \text{or} \quad (2x + 3)^2 + 5(2x + 3) + 6 = 0$$

Before we begin, what do the following equations have in common if you **look at their exponents**?

a. $x^2 + 5x + 6 = 0$

b. $x^{\frac{2}{3}} + 5x^{\frac{1}{3}} + 6 = 0$

c. $x + 5x^{\frac{1}{2}} + 6 = 0$

d. $x^4 + 5x^2 + 6 = 0$

Explain how to solve $x^4 - 13x^2 + 36 = 0$

Examples Solve the following:

a. $x^{\frac{2}{3}} - 2x^{\frac{1}{3}} - 15 = 0$

b. $x - 2x^{\frac{1}{2}} - 8 = 0$

d. $x^{-2} + 8x^{-1} - 9 = 0$

Examples Solve the following:

a. $(x + 3)^2 - 4(x + 3) - 5 = 0$

b. $(2x + 1)^2 - 4(2x + 1) - 21 = 0$

Lesson: Solving Rational Equations

Rational equations are equations that look like:

$$\frac{2x}{x+1} - \frac{3}{x} = \frac{5}{x+1} \quad \text{or} \quad \frac{3}{x^2-4} + \frac{2}{x^2+4x+4} = \frac{5}{x-2}$$

What is the general strategy for solving a rational equation?

What do you have to watch out for with rational equations?

Examples Solve each equation.

a. $\frac{x}{x-4} - 4 = \frac{4}{x-4}$

b. $\frac{3}{x+2} + \frac{10}{x^2-6x-16} = \frac{5}{x-8}$

Examples Solve each equation.

a.
$$-\frac{14}{3x^2+15x-18} = \frac{x}{x-1} + \frac{2}{3x+18}$$

b.
$$\frac{x}{x^2+4x-32} + \frac{2}{x^2-14x+40} = \frac{6}{x^2-2x-80}$$

Lesson: A Review of Solving Radical Equations and Equations with Rational Exponents

What is the general strategy with solving radical equations?

What do you have to watch out for when solving radical equations?

Examples Solve each equation.

a. $x - 3\sqrt{x + 2} = 2$

b. $\sqrt{6x + 7} - 9 = x - 7$

Examples Solve: $\sqrt{3x + 4} - 5 = \sqrt{3x - 11}$

Example Solve each equation.

a. $\sqrt[3]{2x + 1} = \sqrt[3]{3x + 5}$

b. $\sqrt[4]{x + 3} = 2$

Example Solve each equation.

a. $x^{\frac{5}{4}} = 32$

b. $(2x + 1)^{\frac{2}{3}} = 36$

Example Solve for x: $y = k\sqrt{x}$

Lesson: A Review of Solving Equations Quadratic in Form (Using Substitution)

Examples Solve each equation.

a. $x^4 - 10x^2 + 9 = 0$

b. $(x + 1)^{\frac{2}{5}} - 3(x + 1)^{\frac{1}{5}} + 2 = 0$

c. $x^{-\frac{2}{3}} - x^{-\frac{1}{3}} - 6 = 0$

Solving Equations with Radicals

Part 1: One Square Root in the Equation

In this video, we will look at how to solving equations like this:

$$\sqrt{3x+1} = 2 \quad \text{or} \quad \sqrt{4x-2} = 5-x$$

Use the example $\sqrt{3x+1} = 4$ to show the general procedure for solving these types of equations:

Why must we check the solutions?

Example Solve $\sqrt{3x+1} = 8$

Examples Solve

a. $\sqrt{5x - 4} + 2 = 8$

b. $\sqrt{1 + 3x} + 5 = 2$

Explain what is different about solving $\sqrt{5x + 4} = x + 2$

Examples Solve the following.

a. $\sqrt{3x - 6} = x - 2$

b. $\sqrt{2x + 4} = x - 2$

c. $\sqrt{5x + 1} + 5 = 3x$

Solving Equations with Radicals

Part 2: Two Square Roots in the Equation

In this video, we will look at how to solving equations like this:

$$\sqrt{4x - 2} = 3\sqrt{2x + 6} \quad \text{or} \quad \sqrt{3x - 1} + 2 = \sqrt{4x - 2}$$

Let's begin by explaining how to solve: $\sqrt{5x + 2} = \sqrt{3x + 6}$

Examples Solve:

a. $\sqrt{3 - 2x} = \sqrt{2x + 7}$

b. $\sqrt{5 + 3x} = \sqrt{3 + 2x}$

c. $4\sqrt{3 - 2x} = 2\sqrt{1 + 3x}$

Example Solve: $6\sqrt{x} = 4\sqrt{2x+1}$

Now write out the detailed explanation of how to solve $\sqrt{4x+1} - 2 = \sqrt{2x+1}$

Examples Solve:

a. $\sqrt{3x + 6} = 2 + \sqrt{2x - 4}$

b. $\sqrt{3x + 1} = 2 + \sqrt{2x - 2}$

Examples Solve:

a. $\sqrt{2x+4} + \sqrt{1-2x} = 3$

b. $\sqrt{2x+4} = 2 + \sqrt{2x+3}$

Solving Equations with Radicals

Part 3: Two Cube Roots of Higher in the Equation

In this video, we will look at how to solving equations like this:

$$\sqrt[3]{3x+1} = 2 \quad \text{or} \quad \sqrt[4]{2x-1} = 5$$

Explain how to solve: $\sqrt[3]{3x+1} = 2$

Explain how to solve: $\sqrt[4]{3x+4} = 2$

When is it more likely a solution will not work?

Examples Solve the following:

a. $\sqrt[3]{4x + 7} = 3$

b. $\sqrt[3]{3 - 5x} = 2$

c. $\sqrt[4]{3 - 2x} = 3$

d. $\sqrt[4]{4x - 1} = -7$

Rational Expression Applications
Video 1: Proportion Type Problems

What is a proportion?

A bakery can make 300 cookies in 1.5 hours. How long would it take to make 525 cookies?

The ratio of patrons at a café who purchase coffee versus those who purchase tea is 5 to 3. On Monday, the café sold 45 teas. How many coffees were sold?

A painter makes a particular shade of orange by using 3 parts red and 4 parts yellow. The painter has four more gallons of yellow paint than he has of red paint. How many gallons of red paint should he use?

Rational Expression Applications

Video 3: Work Problems

What is the setup for work problems?

It takes Maria 3 hours to paint a room, while it takes her younger sister Lupe 5 hours to paint the same room. How long would it take if the girls painted the room together?

It takes Max 30 minutes to shovel the driveway, while it takes his brother Darrell 45 minutes to do the same job. How long would it take for the guys to do it together?

Rational Expression Applications

Video 2: Distance, Rate, and Time Problems

In these types of problems, we need to use the equation:

$$d = rt$$

Where d =distance, r =rate/speed of a vehicle, and t =time

We can also see that we can rearrange this formula in a few ways:

A boat travels 30 miles downstream in the same time that it takes to travel 20 miles upstream. The current of the river is 5mph. How fast would the boat travel in still water?

A boat travels 16 miles downstream in the same time that it travels 12 miles upstream. In still water, the boat can travel 6mph. What is the speed of the current?

A plane travels 550 miles with the wind. In the same amount of time, it travels 375 miles against the wind. If the wind is blowing at a speed of 25 mph, what is the speed of the plane when there is no wind?

Lesson: A Review of Basic Inequalities

Please check out the examples video on how to graph on a number line, use interval notation, or use set notation in case you forgot!

The Inequality Signs

$<$ Less than

$\leq$ Less than or equal to

$>$ Greater than

$\geq$ Greater than or equal to

The Properties of Inequalities

If a , b , and c are real numbers then:

1. If $a < b$, then $a + c < b + c$

2. If $a < b$ and $c > 0$ then $ac < bc$

3. If $a < b$ and $c < 0$, then $ac > bc$

These properties also hold if you switch to $>$, $\leq$, $\geq$

Examples Solve the following. Put your final answer in interval notation.

$$-2x + 8 < 10$$

Examples Solve the following. Put your final answer in interval notation.

a. $-\frac{1}{3}(x + 3) + \frac{1}{3}x + \frac{2}{5}x \leq \frac{1}{10}$

b. $-5 \leq 1 + 2x < 13$

c. $2 \leq -6x - 3 < 9$

Lesson: Quadratic Inequalities

A quadratic inequality is an inequality in the form: $ax^2 + bx + c < 0$ where a, b, and c are real numbers. You can interchange $<$ with $\leq$, $>$, $\geq$.

There are 3 steps to solving a quadratic inequality. Write out each step as you solve the next example.

Example Write out the steps that you use to solve the following: $x^2 - x - 6 > 0$. Write your answer in interval notation.

Explain Step 1:

Explain Step 2:

Explain Step 3:

Example Solve: $3x^2 - 4 \leq x$. Write your answer in interval notation.

Example Solve $x^2 \geq 25$. Write your answer in interval notation.

Lesson: Rational Inequalities

Rational inequalities look like this:

$$\frac{5}{x+3} > 3 \text{ or } \frac{2x+5}{4x-1} \leq 2$$

We will out the steps to solve a rational inequality by using the following example.

Example Solve $\frac{4}{x+2} \geq 6$ Put your final answer in interval notation.

Explain Step 1 and show how to do it:

Explain Step 2 and show how to do it:

Explain Step 3 and show how to do it:

What do you have to watch out for with rational inequalities?

Example Solve $\frac{3}{x+6} \leq 2$ Put your final answer in interval notation.

Example Solve $\frac{x+3}{x-5} \geq 1$ Put your final answer in interval notation.

Lesson: Number Lines, Interval Notation, and Set Notation

Show how to set up the number line, interval notation, and set notation for the following:

a. $x > 5$

d. $-3 < x < 5$

b. $x \leq 7$

e. $-7 \leq x < -1$

c. $x \geq 0$

f. $3 < x \leq 8$

Lesson: How to Solve Linear Inequalities in One Variable

Linear inequalities in one variable have the form $ax + b < 0$. Alternatively, they could use the symbol $\leq, >, \geq$.

Solving linear inequalities has some similarities and some differences when compared to solving for linear equations.

The Addition and Subtraction Properties

Let a, b, c be real numbers. If $a < b$, then:

This also holds for $\leq, >, \geq$

Examples Solve the following. Put your final answer both on a number line and in interval notation.

a. $x + 4 > 9$

b. $x - 8 \leq 12$

The Multiplication and Division Properties

Let a, b, c be real numbers. If $a < b$, then:

- if c is positive ($c > 0$):

- if c is negative ($c < 0$)

This also holds for $\leq, >, \geq$

Why is this rule different? Use the following example.

To begin, is $-2 < 4$ a true statement? True False

Now multiply both sides of the inequality by 3 and write down the new result below:

This new statement is True False

Now multiply both sides of the inequality by -3 and write down the new result below:

This new statement is True False

Now divide both sides of the inequality by -2 and write down the new result below:

This new statement is True False

Examples Solve the following. Put your final answer both on a number line and in interval notation.

a. $3x \leq 15$

b. $-2x > 8$

c. $5x < -10$

d. $-4x \geq -12$

e. $-9x \leq 27$

From here, you can start to combine the rules to solve more types of inequalities.

Examples Solve the following. Put your final answer both on a number line and in interval notation.

a. $2x - 4 < 6$

b. $3x + 7 > 5x - 9$

c. $\frac{1}{2}x + \frac{1}{5} > \frac{3}{4}x + \frac{7}{10}$

Examples: Solving Linear Inequalities – Basic Examples

Solve the following. Put your final answer both on a number line and in interval notation.

a. $2x + 7 \geq 13$

b. $4x - 6 < 6x - 10$

c. $3(5 - 2x) > 5(3 + 2x)$

d. $2(4 - 2x) - 3(4 - 4x) \leq 4(3x + 3)$

Examples: Solving Linear Inequalities – Examples with Fractions Decimals

Solve the following. Put your final answer both on a number line and in interval notation.

a. $\frac{1}{3}x - \frac{1}{5} \leq \frac{1}{3} + \frac{2}{5}x$

b. $0.03x + 0.2 > 0.02$

c. $\frac{3}{5}(2x - 1) - \frac{3}{4}(2 - 3x) < \frac{1}{10}(4 - 3x)$

d. $0.2(2x - 4) + 0.03(6 - 10x) \geq 0.06(x + 9)$

Lesson: Three-Part Inequalities

Show how to solve the following three-part inequalities. Put your final answer both on a number line and in interval notation.

a. $4 < x + 2 \leq 10$

b. $-3 \leq 2x - 1 < 11$

c. $-5 < 3x + 4 < 19$

d. $4 \leq -1 - 5x < 14$

Lesson: The Basics of Compound Inequalities

The word “and” has special implications in mathematic. One place it is often used is with sets, and we represent “and” with the intersection symbol $\cap$.

The word “or” also has special implications in mathematic. It is also used is with sets, and we represent “or” with the intersection symbol $\cup$.

Examples Given $A = \{1, 2, 3, 4, 5\}$, $B = \{2, 4, 6, 8, 10\}$, $C = \{1, 3, 5, 7\}$, find the following:

a. $A \cap B$

c. $A \cup C$

b. $A \cap C$

d. $B \cup C$

We use similar ideas with compound inequalities.

Show how to represent a solution to $x \geq -2$ and $x < 5$

Example Represent the following with a number line and interval notation.

a. $x > 0$ and $x \leq 7$

b. $x > 8$ and $x \leq -2$

Show how to represent a solution to $x \geq 4$ or $x < -3$

Example Represent the following with a number line and interval notation.

a. $x \leq 0$ or $x > 5$

b. $x > 0$ or $x \leq 5$

We can create more complicated problems where we first have to isolate x , but then we can represent in the same manner as shown above.

Example Represent the following with a number line and interval notation.

$3x - 1 > 5$ and $5x + 4 \leq 24$

Examples: Compound Inequalities

Put your final answer both on a number line and in interval notation.

a. $x - 4 > 8$ or $x + 2 < -3$

b. $2x + 1 \geq 7$ and $5x - 10 < 20$

c. $4x + 2 > 18$ or $3x - 1 < 17$

d. $2 - 3x > 8$ and $5x - 2 > 12$

e. $5x + 9 < 16$ or $4 - 3x > 14$

f. $3 - 2x > 9$ and $5 - x < -12$

Absolute Value Equations

Video 1: The Basics

Absolute value equations look like this:

$$|x + 2| = 7 \quad \text{or} \quad |3x - 5| + 4 = 2$$

To understand how to solve these types of equations, first think about: what are the solutions to $|x| = 5$? Explain why.

From here, we can see that we must consider 2 possible solution sets to find all solutions to absolute value equations. Write down the general strategy for solving absolute value equations:

Solve: $|2x + 3| = 9$

Examples Solve the following:

a. $|3x - 9| = 6$

b. $|7 - 4x| = 3$

To solve $|5x - 4| + 2 = 13$, what must we do first?

Solve:

Examples Solve:

a. $|3x - 6| + 4 = 10$

b. $\left|\frac{1}{2}x + \frac{1}{3}\right| + 5 = 10$

The next video will discuss a special situation and a more complicated equation type.

Absolute Value Equations

Video 2: More Complicated Situations

In this video, we will discuss how to solve equations like:

$$|x + 2| = -7 \quad \text{or} \quad |3x - 5| = |2x + 7|$$

What can we say about the solutions to $|x| = -5$?

Examples Solve the following:

a. $|2x - 7| = -8$

b. $|3x + 5| + 7 = 2$

Absolute value equations can also look like this: $|3x + 1| = |2x - 5|$. Explain how to solve this.

Examples Solve the following:

a. $|5x - 10| = |10x + 15|$

b. $|3x - 5| = |5x + 7|$

c. $|4x - 6| = |3x - 8|$

Example/Question: Why Do You Only Make One Side Negative with Solving $|ax + c| = |bx + d|$ type problems?

What is the approach for solving $|x + 1| = 5$?

What is key to notice here?

How would you solve $|x + 1| = |5x|$?

How would you solve $|x + 1| = |5x - 3|$?

Lesson: Absolute Value Inequalities

Absolute value inequalities have 2 key aspects to solving them. What are those 2 key aspects?

Explain the setup to solve an absolute value inequality of the form $|P| \leq k$ (or $< k$).

Explain the setup to solve an absolute value inequality of the form $|P| \geq k$ (or $> k$).

Let's try to understand why these are setup this way.

Explain the solutions to $|x| < 3$. Summarize what was explained in the video.

Explain the solutions to $|x| \geq 3$. Summarize what was explained in the video.

Examples Solve the following inequalities. Graph your solution set on a number line and express your solution in interval notation.

a. $|2x - 4| < 6$

b. $|3x + 6| \geq 9$

c. $|x - 5| - 4 \leq 6$

d. $5 \geq |3x - 4|$

e. $7 < 3 + |x - 4|$

f. $6 \leq 4 + |8 - 2x|$

Lesson: Special Absolute Value Inequalities

We will discuss some special situations. Explain the reasoning for each of these situations:

a. $|2x + 3| + 7 < 4$

b. $|3x - 2| > -3$

c. $|5x + 1| \geq 0$

d. $|2x - 4| < 0$

e. $|6x + 2| \leq 0$

Lesson: A Review of Absolute Value Equations and Inequalities

We can summarize how to solve each case with the following chart:

Case (k is a number)	How to Setup
$ x = k$	
$ x < k$ (or $\leq k$)	
$ x > k$ (or $\geq k$)	

Examples Solve the following. Put your answer on a number line and in interval notation when you have an inequality.

a. $|x + 3| = 5$

b. $|2x - 4| \geq 6$

Examples Solve the following. Put your answer on a number line and in interval notation when you have an inequality.

a. $|4x - 2| = |6x + 8|$

b. $|5x - 2| + 1 < 9$

c. $|2 - x| - 1 > 9$

Special Cases!

Example Explain what is special about each case below and how to solve the problem.

a. $|4x - 1| = -7$

b. $|4x - 1| = 0$

c. $|4x - 1| < -7$

d. $|4x - 1| > -7$

Example Explain what is special about each case below and how to solve the problem.

a. $|4x - 1| \geq 0$

b. $|4x - 1| \leq 0$

c. $|4x - 1| < 0$

d. $|4x - 1| > 0$

Lesson: An Intro to Functions

Define a relation:

Define a domain:

Define a range:

Example In the set of ordered pairs $\{(2,3), (4, -1), (7, 2)\}$, define the domain and range of the relation.

Define a function:

What is the visual explanation that explains what a function is or is not?

Example In the set of ordered pairs $\{(2,3), (4, -1), (7, 2)\}$, is this a function?

Example Are the following functions?

a. $\{(4, -1), (3, 2), (7, -1)\}$

b. $\{(2, 4), (3, 5), (2, -1)\}$

What is the notation for a function? How do we describe/use this notation?

Example In the set of ordered pairs $\{(2, 3), (4, -1), (7, 2)\}$, what is $f(4)$?

Example Suppose $f(x) = 2x + 1$ and $g(x) = x^2 - 3$. Find the value of the following:

a. $f(2)$

b. $g(1)$


c. $f(a)$

d. $g(h + k)$


What is the vertical line test?

Example Do the following graphs represent a function?

a.



b.




Lesson: Determining the Domain and Range of Functions In General

How do you determine the domain and range of a set of points like $\{(2,4), (5,8), (4,6)\}$?

What is the domain and range of that set of points?

In general, it is easiest to determine the domain and range when we have the graph in front of us.

For the following graph, write out how you should determine the domain and range.



Write a sentence on how you should find the domain:


State the domain:

Write a sentence on how you should find the range:

State the range:

Examples Determine if the following graphs represent functions, and then state the domain and range.


a.



Domain:

Range:


b.



Domain:

Range:

c.



Domain:

Range

Lesson: The Domain of Some Specific Types of Functions

What is a polynomial? Write some examples.

What is the domain of any polynomial function?

What is a rational function?

What is the domain of any rational function?

Examples Determine the domain of the following.

a. $f(x) = 3x - 1$

b. $g(x) = \frac{4}{x+2}$

c. $h(x) = 3x^3 - 2x^2 + 4x - 1$

d. $f(x) = \frac{2x+1}{x^2-9}$

What is a square root function?

What is the domain of a square root function?

Examples Find the domain of the following:

a. $f(x) = \sqrt{x - 2}$

b. $g(x) = \sqrt{4x - 1}$


c. $h(x) = \sqrt{4 - x}$

Lesson: How to Graph a Function


How do you graph a function?

Examples Graph the following functions.

a. $f(x) = 3x + 1$



b. $g(x) = 4$



Lesson: Review of Functions

How do you represent a function?

What is a function often interchangeable with? How can you rewrite $f(x) = 3x + 8$?

So what IS a function? What does it actually represent?

What's another handy feature of functions?

What are the domain and range?

Does domain and range only work for functions?

Examples Are the following functions, and what is the domain and range?


a. $\{(2,3), (5,-1), (-3,7)\}$

b. $\{(-1,5), (-8,3), (-1,6)\}$


What is the Vertical Line Test, and how does it help you determine if you have a function?

Example State the domain and range of the following, and use the vertical line test to determine if the graph represents a function.

a.



b.



Examples Determine if the following represent functions.

a. $y = 3x^2$

b. $x = y^2$

c. $y = \frac{5}{x^2}$

Example Write the following in function notation: $3x + 2y = 6$

Example Suppose $f(x) = 3x - 5$ and $g(x) = x^2 + 4x - 1$ and $h(x) = \{(2, 5), (6, -3), (-1, -4)\}$. Find the following:

a. $f(7) =$

b. $g(3) =$

c. $f(k) =$

d. $g(x + h) =$

e. $h(6) =$

Lesson: Increasing, Decreasing, and Constant Functions

Below are the “formal math” definitions of whether a function is increasing, decreasing, or constant. Let I be an open interval and let x_1 and x_2 be two points in that interval.

- A function is **increasing** over I if when $x_1 < x_2$, $f(x)_1 < f(x_2)$
- A function is **decreasing** over I if when $x_1 < x_2$, $f(x)_1 > f(x_2)$
- A function is **constant** over I if when $x_1 < x_2$, $f(x)_1 = f(x_2)$

While these are the formal definitions, what we really want to know are the intuitive meanings of these definition.

Sketch a graph that is increasing:

Decreasing:

And constant:

Example Sketch the graph in the video, and then identify where the graph is increasing, decreasing, or constant.

a.

b.

Lesson: A Few Basic Graphs

There are a few graphs that are helpful to immediately know as we move forward. Let's break these down.

The Parabola: $f(x) = x^2$

Create table of relevant points and then sketch your own graph.

Absolute Value: $f(x) = |x|$

Create table of relevant points and then sketch your own graph.

X cubed: $f(x) = x^3$

Create table of relevant points and then sketch your own graph.

Square Root: $f(x) = \sqrt{x}$

Create table of relevant points and then sketch your own graph.

Cube Root: $f(x) = \sqrt[3]{x}$


Create table of relevant points and then sketch your own graph.

Lesson: An Intro to Translations and Reflections of Graphs

We are going to develop the intuition for this concept before we actually graph anything.

Vertical Translations

Draw the base graph of $f(x) = x^2$ in one color and then use some other colors to show how the translations work. Label the graphs.




Explain the general way that vertical shifts work:

Example How does $f(x) = \sqrt{x} + 6$ shift?

Horizontal Translations

Draw the base graph of $f(x) = |x|$ in one color and then use some other colors to show how the translations work. Label the graphs.




Explain the general way that horizontal shifts work:

Example How does $f(x) = \sqrt[3]{x+3}$ shift?

Reflections Over the X-Axis

Draw the base graph of $f(x) = \sqrt{x}$ and then graph the reflection in another color. Label the graph of the reflection. Label the graphs.



Explain the general way that reflections over the x-axis work:

Example How does $f(x) = -x^2$ look?

Example Describe the translations and reflections of the following.

a. $f(x) = -x^2 + 3$

b. $g(x) = \sqrt{x+2} - 1$

c. $h(x) = (x-5)^3 + 2$

d. $f(x) = -|x+2| + 5$


e. $g(x) = -\sqrt[3]{x-1} - 4$

Lesson: Graphing with Translations and Reflections

Examples For each of the following, describe the translations and reflections. Then graph with 2-3 points.


a. $f(x) = x^2 + 1$

Describe any translations or reflections:




b. $g(x) = \sqrt{x - 2}$

Describe any translations or reflections:




c. $f(x) = (x + 2)^3 - 1$

Describe any translations or reflections:




d. $h(x) = |x - 3| + 2$

Describe any translations or reflections:




e. $f(x) = -|x - 3| - 2$

Describe any translations or reflections:



f. $f(x) = -\sqrt{x + 3} + 1$

Describe any translations or reflections:



Lesson: Writing Equations of Functions with Translations and Reflections

Example Write the equation of a function $g(x)$ where the following transformations are performed on the graph of $f(x)$

a. $f(x) = \sqrt[3]{x}$ is shifted 3 units down and 2 units right

b. $f(x) = x^2$ is shifted 2.5 units up and π units left

c. $f(x) = e^x$ is reflected about the x-axis, shifted 4 units down and 1 unit left

Lesson: Graphs of Basic Functions

In this section, we will review some basic graphs that are vital to know for this class and calculus. Knowing the “basic shapes” of certain graphs will make your life much easier and give you greater intuition when graphing.

Show how to graph the following and highlight the **key points** and **key patterns**. This will come in handy for the next lesson. Take your time to sketch your own NEAT CLEAN CLEAR graphs.

$$f(x) = x$$

$$f(x) = x^2$$

$$f(x) = x^3$$

$$f(x) = \sqrt{x}$$

$$f(x) = \sqrt[3]{x}$$

$$f(x) = |x|$$

Lesson: Piecewise Functions

What is a piecewise function? Draw an example of what they look like.

Example Given the function: $f(x) = \begin{cases} 3x - 1, & x \geq 2 \\ \frac{1}{2}x - 4, & x < 2 \end{cases}$ evaluate the following:

a. $f(3) =$

b. $f(-1) =$

c. $f(2) =$

Example Given the function: $f(x) = \begin{cases} 4, & x > 3 \\ 4x - 2, & -2 < x \leq 3 \\ x^2 - 5, & x \leq -2 \end{cases}$ evaluate the following:

a. $f(4) =$


b. $f(-5) =$

c. $f(-2) =$


d. $f(3) =$

Example Show how to graph the following piecewise functions.


a. $f(x) = \begin{cases} 4, & x > 5 \\ -1, & x \leq 5 \end{cases}$




b. $f(x) = \begin{cases} 3x + 1, & x > -3 \\ -2x, & x \leq -3 \end{cases}$



$$c. f(x) = \begin{cases} 3, & x > 2 \\ 2x, & -1 < x \leq 2 \\ -x + 3, & x \leq -1 \end{cases}$$



$$d. f(x) = \begin{cases} 3x + 1, & x \geq 1 \\ 6, & -1 \leq x < 1 \\ x^2, & x < -1 \end{cases}$$



Lesson: The Greatest Integer Function

Show how to graph the greatest integer function $f(x) = \llbracket x \rrbracket$

Example Graph $f(x) = \left\llbracket \frac{1}{2}x - 2 \right\rrbracket$

Lesson: Review of Function Operations

Given that f and g are functions, how do we define the function operations?

$$(f + g)(x) =$$

$$(f - g)(x) =$$

$$(fg)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

Example Let $f(x) = 3x + 1$, $g(x) = x^2 - 4$, $h(x) = 3x^2 + 7x + 2$, and $k(x) = x^2 + 6x + 9$. Find each of the following:

a. $(f + g)(x) =$

b. $(f + g)(3) =$

c. $(g - k)(1) =$

Example Let $f(x) = 3x + 1$, $g(x) = x^2 - 4$, $h(x) = 3x^2 + 7x + 2$, and $k(x) = x^2 + 6x + 9$. Find each of the following:

a. $\left(\frac{h}{f}\right)(x) =$

b. $(gh)(x) =$

c. $\left(\frac{k}{h}\right)(1) =$

Lesson: Review of Composition of Functions

For two functions f and g , what is the composition $(f \circ g)(x)$?

Example Let $f(x) = 3x + 1$, $g(x) = x^2 - 4$, $h(x) = \sqrt{x}$, and $k(x) = x^2 + 6x + 9$. Find each of the following:

a. $(f \circ g)(x) =$

b. $(g \circ f)(x) =$

Example Let $f(x) = 3x + 1$, $g(x) = x^2 - 4$, $h(x) = \sqrt{x}$, and $k(x) = x^2 + 6x + 9$. Find each of the following:

a. $(k \circ f)(1) =$

b. $(g \circ h)(4) =$

c. $(h \circ k)(x) =$

Lesson: The Difference Quotient

Explain what the difference quotient is and draw some sketches to explain how it works. In other words, watch the video explanation and then explain it in your own words on this page.

State the formula for the difference quotient here:

Example Calculate the different quotient for the following.

a. $f(x) = 4x + 5$

b. $f(x) = x^2 - 1$

Lesson: The Difference Quotient

Find the difference quotient.

a. $f(x) = \frac{1}{x+3}$

b. $f(x) = x^2 - 4x + 1$

c. $f(x) = -x^2 + 3$

Lesson: Translations and Reflections of Functions

Using the graph $f(x) = |x|$ as the “base function,” sketch how vertical translations work (summarize what was discussed in the video).

How do you describe vertical translations in function notation?

Using the graph $f(x) = x^2$ as the “base function,” sketch how horizontal translations work (summarize what was discussed in the video).

How do you describe horizontal translations in function notation?

Use the graph $f(x) = \sqrt{x}$ to show how reflections over the x- and y-axis work (summarize what was discussed in the video).


How do you describe reflections over the x-axis and y-axis using function notation.

Use the graph $f(x) = |x|$ to show how stretching and shrinking works (summarize what was discussed in the video).


Explain how stretching and shrinking works with function notation.

Example Describe any translations / reflections / stretching / shrinking and sketch a graph of each of the following.

a. $f(x) = -x^2 + 3$




b. $g(x) = (x + 2)^3 - 3$




Example Describe any translations / reflections / stretching / shrinking and sketch a graph of each of the following.

a. $g(x) = -2(x + 1)^2 - 5$



b. $h(x) = 2\sqrt{-x} + 1$



Examples Write the equation of the given functions with the following characteristics.

a. The function $f(x) = x^2$ moved 3 units left and 5 units down.

b. The function $g(x) = \sqrt[3]{x}$ reflected over the y-axis, translated 2.5 units right and 1.75 units up.


Transformations of Functions

These are looking at graphs of things like:


$$f(x) = x^2 - 5 \quad \text{or} \quad f(x) = |x + 3| - 2 \quad \text{or} \quad f(x) = \sqrt{x - 3}$$

First we need to discuss the shape of a few basic functions. Write these down and note the important points.


a. $f(x) = x^2$



b. $f(x) = |x|$




c. $f(x) = \sqrt{x}$



Using a few different colors, sketch out what the graphs of

a. $f(x) = x^2$ versus b. $g(x) = x^2 + 3$ versus c. $h(x) = x^2 - 1$

look like all on the same graph below:




This is an example of a vertical shift. Explain what this means in mathematical terms:

Using a few different colors, sketch out what the graphs of

a. $f(x) = |x|$ versus b. $g(x) = |x - 1|$ versus c. $h(x) = |x + 2|$

look like all on the same graph below:




This is an example of a horizontal shift. Explain what this means in mathematical terms:

Using a few different colors, sketch out what the graphs of

a. $f(x) = |x|$ versus b. $g(x) = -|x|$


look like all on the same graph below:



This is an example of a reflection over the x-axis. Explain what this means in mathematical terms:


Example Explain in plain language what types of shifts/reflections are occurring, and then graph the function.

$$f(x) = |x| + 3$$




Examples Explain in plain language what types of shifts/reflections are occurring, and then graph the function.


a. $f(x) = (x - 1)^2$




b. $g(x) = \sqrt{x} - 2$




c. $h(x) = |x + 1| - 2$




d. $k(x) = (x - 2)^2 + 4$




e. $f(x) = -(x + 3)^2 - 2$




f. $g(x) = -|x + 3| - 1$



g. $h(x) = -\sqrt{x - 2}$



h. $k(x) = -\sqrt{x + 1} - 3$



Examples Write the equations that represent the listed transformations.

a. A function $f(x) = \sqrt{x}$ is shifted 7 units down.

b. A function $g(x) = x^3$ is shifted 3 units left and 2 units up.

c. A function $h(x) = |x|$ is shifted 2.5 units right and reflected over the x-axis.


d. A function $f(x) = e^x$ is shifted 4 units up and reflected over the x-axis.

e. A function $g(x) = \sqrt[3]{x}$ is shifted 3.2 units right, 5.7 units up, and reflected over the x-axis


Examples: Translations and Reflections of Functions

Describe any translations / reflections / stretching / shrinking and sketch a graph of each of the following.


a. $f(x) = -(x + 2)^2 - 3$



b. $g(x) = \frac{1}{2}(x - 1)^3 + 2$



c. $h(x) = -\frac{1}{3}\sqrt{x + 1} + 1$



Lesson: Symmetry and Even or Odd Functions

Explain when a graph is symmetric with respect to the y-axis:

Explain when a graph is symmetric with respect to the x-axis:

Explain when a graph is symmetric with respect to the origin:

What makes a function odd?

What makes a function even?

So what is the test for even or odd functions?

Examples Determine what type of symmetry the following functions have:

a. $y = x^2 + 4$

b. $x^2 + y^2 = 25$

Examples Determine what type of symmetry the following functions have:

a. $y = x^3 - 5$

b. $y = 7x^2 + 2x - 4$

c. $y = 2x^{\frac{1}{3}} - 3x$

Lesson: Determining if Functions are One-to-One

Before we begin, you should already be familiar with the concept of functions and the vertical line test. First let's do some review.

Review Determine whether the following relations are functions, then determine the domain and range.

a. $\{(2, 3), (5, 7), (2, 4)\}$

b.

Now we will discuss a specific type of function: a one-to-one function. Define this below:

How can we visualize this?

Example Determine if any of the following are i) functions and ii) one-to-one functions

a. $\{(2, 4), (-1, 7), (3, 7)\}$

b. $\{(3, 6), (-1, 2), (-3, -4)\}$

Similar to how we have a test to determine if a graph represents a function, we have another test to determine if a function is one-to-one.

Describe the horizontal line test below:

Examples Determine whether the following graphs represent i) functions and ii) one-to-one functions (draw your own):

a.

b.

c.

Lesson: Finding the Inverse of Functions


One-to-one functions have inverses. Describe what the inverse of a function is:

Example Find the inverse of the following function: $\{(1, 4), (7, 8), (3, -2)\}$


Describe the steps to find the inverse of a function:

Examples Find the inverse of the following one-to-one functions. Then plot the original function on the graph with its inverse.

a. $f(x) = 3x + 1$



b. $g(x) = -\frac{1}{2}x + 4$



What is the relationship of a graph with its inverse?

Examples Find the inverse of the following one-to-one functions.

a. $h(x) = \sqrt[3]{x}$

b. $f(x) = x^2, x \geq 0$

Lesson: A Review of Inverse Functions

What is a one-to-one function?



Examples Are the following one-to-one functions? If no, why not?

a. $f(x) = x^2$

b. $g(x) = 2x + 1$

What is the horizontal line test?

Example Use the horizontal line test to determine if the functions are one-to-one:



What is an inverse function?

What does an inverse function have to do with one-to-one functions?

Describe how to use composition to determine if 2 functions are inverses of one another.

Example Use composition to determine if the functions $f(x) = 2x - 4$ and $g(x) = \frac{1}{2}x - 2$ are inverses.

Example Use composition to determine if the functions $f(x) = \frac{2}{x+6}$ and $g(x) = \frac{6+x}{2}$ are inverses.

Using $f(x) = 4x + 2$, outline the steps for how to find the inverse of a function.

Example Find the inverse of the following functions.

a. $f(x) = \sqrt[3]{3x + 1}$


b. $f(x) = \frac{x+2}{x-1}$

c. $f(x) = \sqrt{x - 5}$


How can we tell on a graph if 2 functions are inverses of one another?

Examples Are the following graphs inverses of one another?


a.



b.



Example Draw the inverse of the following graph.



Lesson: Review of Lines and Linear Functions

What makes a function **linear**? What are 2 forms of a linear function? How do you interpret this equation?

Examples What is the domain and range of $f(x) = 4x + 1$. Show how to graph it.

Examples What is the domain and range of $f(x) = 5$. Show how to graph it.

Examples What is the domain and range of $x = -4$. Show how to graph it.

What is the **standard form** of a line?

Examples Give the domain and range of $3x - y = 6$. Show how to graph it.

How do you find the **slope** of a line?

What is the slope of a vertical line?

What is the slope of a horizontal line?

Examples Find the slopes between the following sets of points.

a. $(5, 6)$ and $(-3, 2)$

b. $(-4, 7)$ and $(2, 7)$

c. $(6, 2)$ and $(6, -1)$

Example Find the slope of the following lines:

a. $4x - 2y = 8$

b. $-5x + 3y = -15$

Lesson: Review of Writing Equations of Lines

What is the point-slope form of a line?


Example Write the equation of a line through the point $(4, 3)$ with slope $m = -2$.

Example Write the equation of a line that goes through the points $(2, 4)$ and $(-4, 10)$. Put your answer in standard form.

What is the slope-intercept form of a line?

Example Show how to write $4x - 6y = 12$ in slope-intercept form.

Example Find the slope, x- and y-intercept, and equation of the line:




What is the form of a vertical line?


What is the form of a horizontal line?

Example Determine the equation of the following lines.

a.



b.



What makes 2 lines parallel?

What makes 2 lines perpendicular?

Examples Write the equation of a line that goes through $(-2, -5)$ that satisfies the conditions:

a. parallel to the line: $3x - 2y = -6$

b. perpendicular to the line: $x - 2y = -4$

Quick Reference of Line Formulas

Slope-Intercept Form

Standard Form

Point-Slope Form

Slope Formula

Form of a Horizontal Line

Form of a Vertical Line

What makes 2 lines parallel?

What makes 2 lines perpendicular?

Examples: Writing the Equation of a Line Through 2 Points

Example Write the equation of a line through the given 2 points. Put your answer in standard form, and your standard form should not include any fractions or decimals.

a. $(4, 5)$ and $(2, 9)$

b. $(2, 7)$ and $(-4, 7)$

c. $(-4, 2)$ and $(2, 5)$

d. $(3, 5)$ and $(3, 8)$

Examples: Writing the Equation of a Line Through 2 Points

Example Write the equation of a line through the point $(2, 3)$ that satisfies the given conditions. Your final answer should be in standard form with no fractions or decimals.

a. Parallel to $5x - 2y = 10$

b. Perpendicular to $4x - 2y = 1$

c. Parallel to $y = 5$

d. Perpendicular to $x = 7$

Lesson: The Slope of a Line


What does slope represent?

Draw the various possible slopes of lines:


How do find the slope of a line?

Examples Find the slope of the following lines:

a.



b.



What is the slope formula?

Examples Use the slope formula to find the slope between the pairs of points:

a. $(3, 5)$ and $(4, 8)$

c. $(7, 2)$ and $(7, 4)$

b. $(6, -2)$ and $(-2, 4)$


d. $(2, 4)$ and $(-1, 4)$

Lesson: Using Slope to Graph Lines; Intro to Slope-Intercept Form


When the slope of a line and a point are known, a line can be easily graphed.

Examples Given the slope and the point, graph the following lines:


a. $m = 2$ and $(-1, 2)$




c. $m = 1$ and $(2, -1)$



b. $m = -\frac{1}{3}$ and $(-3, 4)$



d. $m = \frac{2}{3}$ and $(0, -3)$




What is the slope-intercept form of a line?

Examples Identify the slope m and the y-intercept $(0, b)$ in each of the following equations:


Equation	Slope, m	What stands for b	State the y-intercept (as a point)
a. $y = -2x + 3$			
b. $y = \frac{1}{4}x - 7$			
c. $y = 3x + 9$			
d. $y = x$			

Examples Graph the following lines:


a. $y = 2x + 3$




c. $y = \frac{1}{3}x - 4$



b. $y = -x + 4$



d. $y = 3x$



Lesson: Rewriting Linear Equations into Slope-Intercept Form of a Line

What is the slope-intercept form of a line?

Explain how to put the equation $2x - 4y = 8$ into slope-intercept form.

Examples Put the following equations in slope-intercept form:

a. $6y - 3x = 12$

b. $-5x + 2y = -4$

Lesson: Determining Whether Two Lines are Parallel or Perpendicular

What makes 2 lines parallel?

Examples Are the equations $y = \frac{2}{3}x + 5$ and $2x - 3y = 6$ parallel?

What makes 2 lines perpendicular?

Examples Determine the negative reciprocal of each of the following:

a. $\frac{2}{3}$

c. -7

b. $-\frac{1}{5}$

d. 1

Example Are the equations $y = \frac{3}{5}x + 2$ and $-3x + 5y = -15$ perpendicular?

Examples Are the following pairs of lines parallel, perpendicular, or neither?

a. $3x - 2y = 6$ and $4y = -6x - 8$

b. $y - 3x = 0$ and $12x - 4y = 24$

c. $y = 3$ and $x = -2$

Examples Lines A and B contain the given points. Determine whether the lines are parallel, perpendicular, or neither.

a. Line A: $(2, 11), (5, 10)$ Line B: $(1, 3), (-6, -18)$

b. Line A: $(-5, 5), (-7, -2)$ Line B: $(0, 11), (-4, 9)$

Lesson: Putting Equations in Standard Form

What is the standard form of a line?

Examples Write the following equations in standard form:

a. $y = 2x - 4$

b. $y = -5x + 3$

c. $y = -\frac{1}{2}x + 1$

d. $y = \frac{2}{3}x - 2$

Lesson: Writing Equations of Lines with The Point-Slope Form

What is the point-slope form of a line?

Examples Given the point and the slope, write the point-slope form of a line.

a. $m = -2, (1, 3)$

b. $m = -\frac{2}{3}, (5, -2)$

What is the slope-intercept form of a line?

What is the standard form of a line?

Examples Given the point and the slope, write the equation of the line in the desired format.

a. $m = -1$, $(2, -5)$ in slope-intercept form

b. $m = 3$, $(5, -4)$ in standard form

c. $m = -\frac{1}{2}$, $(6, 3)$

d. $m = \frac{2}{3}$, $(-5, -1)$

Lesson: Writing the Equation of a Line from Two Points

Write the equation of the line that contains the following 2 points in the desired format.

a. $(4, 2)$ and $(5, 5)$ in slope-intercept form

b. $(-2, 4)$ and $(-4, -2)$ in standard form

c. $(5, -3)$ and $(3, 5)$ in slope-intercept form

d. $(-1, -3)$ and $(1, 5)$ in standard form

Lesson: Writing the Equation of a Line Parallel/Perpendicular to Another Line

What makes 2 lines parallel?

What makes 2 lines perpendicular?

Examples Write the equation of a line with the given conditions.

a. Parallel to $y = 3x + 1$ that goes through $(-2, -5)$ in slope-intercept form

b. Perpendicular to $y = -\frac{2}{3}x - 5$ that goes through $(6, -2)$ in standard form

c. Parallel to $2x - 3y = 6$ that goes through $(1, -2)$ in slope-intercept form

Examples Write the equation of a line with the given conditions.

a. Perpendicular to $2x - 4y = 8$ that goes through $(-4, -2)$ in standard form

b. Parallel to $3x - y = 12$ that goes through $(-1, -5)$ in slope-intercept form


c. Perpendicular to $-2x + 3y = 12$ that goes through $(2, -5)$ in standard form

Lesson: Understanding the Graphs of Quadratic Functions


Quadratic functions are functions of the form _____, where $a \neq 0$. But in previous lessons we worked on graphing quadratic functions, ie parabolas in another way. Let's review that first.

Examples Graph the following:

a. $f(x) = (x + 2)^2 - 3$




b. $g(x) = -(x - 3)^2 + 1$



To begin, we will add on one more way to manipulate the graph of a parabola. Explain how to graph the follow:


a. $f(x) = 3x^2$

How do you interpret the 3?




b. $g(x) = -\frac{1}{2}(x - 2)^2 + 1$

How do you interpret the $-\frac{1}{2}$?




Examples Graph the following:

a. $f(x) = -2(x + 1)^2 + 3$



b. $g(x) = \frac{1}{3}(x - 2)^2 - 2$




Using all of the examples we have discussed, what is another equation form for the parabola?

Explain some of the characteristics of a parabola that we get from this format:

Examples For each of the following functions, find and state: (i) the vertex (ii) axis of symmetry (iii) x-intercepts (iv) y-intercepts (v) and then graph the function

a. $f(x) = (x - 3)^2 - 4$




Vertex:

Axis of Symmetry:

X-intercepts:

Y-intercepts:

b. $f(x) = -2(x - 3)^2 + 8$



Vertex:


Axis of Symmetry:

X-intercepts:

Y-intercepts:

Examples For each of the following functions, find and state: (i) the vertex (ii) axis of symmetry (iii) x-intercepts (iv) y-intercepts (v) and then graph the function

a. $f(x) = \frac{1}{4}(x + 2)^2 - 3$




Vertex:

Axis of Symmetry:

X-intercepts:

Y-intercepts:

b. $f(x) = -\frac{1}{3}(x - 1)^2 + 3$



Vertex:

Axis of Symmetry:


X-intercepts:

Y-intercepts:

Examples: Understanding How Many X-Intercepts There Are

Examples For each of the following functions, find and state: (i) the vertex (ii) axis of symmetry (iii) x-intercepts (iv) y-intercepts (v) and then graph the function

a. $f(x) = \frac{1}{2}(x + 1)^2 - 8$




Vertex:

Axis of Symmetry:

X-intercepts:

Y-intercepts:

b. $f(x) = -2(x - 3)^2$



Vertex:


Axis of Symmetry:

X-intercepts:

Y-intercepts:

Examples For each of the following functions, find and state: (i) the vertex (ii) axis of symmetry (iii) x-intercepts (iv) y-intercepts (v) and then graph the function

a. $f(x) = (x - 2)^2 + 1$




Vertex:

Axis of Symmetry:

X-intercepts:

Y-intercepts:

b. $f(x) = -(x + 3)^2 - 1$



Vertex:

Axis of Symmetry:

X-intercepts:

Y-intercepts:

Lesson: Finding the Vertex of Quadratic Equations in the form $f(x) = ax^2 + bx + c$

What is the vertex formula?

Show how to use the vertex formula for $f(x) = 3x^2 + 6x - 2$


Examples Find the vertex of the following:

a. $f(x) = -x^2 + 4x - 2$

b. $f(x) = -\frac{1}{2}x^2 + 3x - 1$

Lesson: Putting Quadratics into the $f(x) = a(x - h)^2 + k$ Form

Before you begin this lesson, you should already feel comfortable graphing an equation like $f(x) = 2(x - 1)^2 + 3$. Graph this below:



Now show all the steps to put $f(x) = 2x^2 - 8x + 2$ into $f(x) = a(x - h)^2 + k$ form:

Examples Put the following quadratic functions into $f(x) = a(x - h)^2 + k$ form:

a. $f(x) = x^2 + 4x - 2$

b. $g(x) = x^2 + 6x + 10$

c. $h(x) = -x^2 + 2x - 4$

d. $y = 3x^2 + 6x - 12$

Examples: Putting Quadratics into the $f(x) = a(x - h)^2 + k$ Form

Examples Put the following quadratic functions into $f(x) = a(x - h)^2 + k$ form:

a. $f(x) = \frac{1}{2}x^2 + x - 3$

b. $g(x) = -\frac{1}{3}x^2 + 4x - 2$

c. $y = x^2 + 3x + 2$

Lesson: Determining Whether a Quadratic Function will have a Minimum or a Maximum

Using the form $f(x) = ax^2 + bx + c$, explain what determines whether the parabola opens up or down:

Examples Determine whether the following quadratics open upwards or downwards, and then determine their vertex:

a. $f(x) = -2x^2 + 4x - 2$ upwards downwards

b. $g(x) = -\frac{1}{2}x^2 - 2x + 2$ upwards downwards

c. $h(x) = x^2 + 6x + 1$ upwards downwards

Lesson: Applications of Quadratic Functions 1

A ball is thrown upwards from a cliff 24 feet about the ground. The height h of the ball, in feet, t seconds after it is thrown is given by:

$$h(t) = -16t^2 + 32t + 24$$

How long does it take the ball to reach its maximum height?

What is the maximum height attained by the ball?

Lesson: Review of Quadratic Functions

A polynomial function of degree n is a function of the form

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

Where $a_n, a_{n-1}, \dots, a_1, a_0$ are any complex or real number, $a_n \neq 0$, and n is a nonnegative integer.

What do we call n in this equation?

What do we call a_n in this equation?

What do we call $a_n x^n$ in this equation?

Example What is the degree, leading coefficient, and dominating term in the following?

a. $f(x) = 4x^4 - 5x^2 + x^7 + 100$

b. $g(x) = x\sqrt{3} - 5$


c. $h(x) = 7$

What is a quadratic function?

What do we call the graph of a quadratic function?

Draw and label the parts of a parabola.

Example Sketch the graph of $f(x) = -(x + 4)^2 - 2$



What is the **vertex form** of the equation of the parabola?

Example Show how to use completing the square to put $f(x) = x^2 + 6x - 4$ in vertex form. Write out clear directions.

Example Use completing the square to write $f(x) = 3x^2 + 5x + 2$ in vertex form, then sketch the graph.

Example Find the intercepts of $f(x) = 3x^2 + 5x + 2$

How many x-intercepts can a parabola have? Draw a little sketch of each situation.

How can you tell how many intercepts a parabola will have if the parabola is in the form: $f(x) = ax^2 + bx + c$

What is the vertex formula for a parabola in the form $f(x) = ax^2 + bx + c$?

Example Use the vertex formula to find the vertex of the following: $f(x) = 2x^2 - 4x + 1$

Examples: Finding Intercepts, Vertices, Etc of Quadratic Functions

Find the vertex, axis of symmetry, domain, range, intercepts, and intervals of increasing and decreasing. If the function is in $f(x) = ax^2 + bx + c$ form, put it in vertex form. Sketch the graph. If the function is in $f(x) = a(x - h)^2 + k$ form, put it into standard form.

a. $f(x) = -\frac{1}{2}x^2 - 3x - \frac{1}{2}$

b. $f(x) = 2(x - 1)^2 + 3$

Lesson: Graphs of Basic Polynomial Functions and Translations of the Basic Graphs

Explain the graphs of $f(x) = x^2$, $f(x) = x^4$, $f(x) = x^6$, etc.

Explain the graphs of $f(x) = x^3$, $f(x) = x^5$, $f(x) = x^7$, etc.

If we multiply the polynomial like so: $f(x) = ax^n$, what happens to the graph?

What happens if we have $f(x) = a(x - h)^n + k$?

Example Sketch the graphs:


a. $f(x) = (x - 5)^5 + 3$

b. $f(x) = -2(x + 1)^6 - 4$


Lesson: Graphs of Polynomial Functions, Turning Points, and End Behavior

In the last video, we discuss the graphs of basic polynomials and translations. In this video, we will look at more complicated possibilities.

Below are the graphs of a few polynomials. What patterns do you notice?




$$f(x) = x^3 - 2x^2 - 5x + 6 = (x - 1)(x + 2)(x - 3)$$




$$\begin{aligned} f(x) &= x^5 - 7x^4 + 19x^3 - 37x^2 + 60x - 36 \\ &= (x - 3)^2(x - 1)(x - 2i)(x + 2i) \end{aligned}$$

$$f(x) = (x - 2)^2(x + 1)^3(x + 2)$$



$$f(x) = (x - 1)^2(x + 1)(x - 2)$$



What do you notice about the behavior of these graphs at zeros? Summarize this in your own words, it was discussed as a general pattern in the video.

What are turning points and how many will a degree n polynomial have?

What is the end behavior of a polynomial and how do you determine it?

Example What are the number of turning points of the following and what is the end behavior of each?

a. $f(x) = x^2 + 3x + 2$

b. $f(x) = (x + 1)^2(x - 2)$

c. $f(x) = -\frac{1}{2}x(x - 3)^4(x + 1)^3(x + 2)$

List the steps required to graph a polynomial:

Example Graph $f(x) = x^3 - x^2 - 4x + 4$

Example Graph $f(x) = -(x + 1)(x - 2)^2(x - 3)$

What is the relationship between x-intercepts, zeros, solutions, and factors of polynomials???

Examples: Graphs of Polynomial Functions, Turning Points, and End Behavior

Graph the following polynomials:

a. $f(x) = x^3 + 5x^2 + 2x - 8$

b. $f(x) = 3x^4 + 5x^3 - 2x^2$

c. $f(x) = (x - 1)^3(x + 2)^2$

Lesson: The Intermediate Value Theorem and the Boundedness Theorem

State the Intermediate Value Theorem

Example Using synthetic division to show $f(x) = -x^3 + 2x^2 - 4x + 5$ has a real zero between 1 and 2.

Example Using synthetic division to show $f(x) = x^4 - 4x^3 - x + 3$ has a real zero between 0.5 and 1.

State the Boundedness Theorem

Example Show that $f(x) = x^4 + 5x^2 + 3x - 7$ has no real zero less than -3.

Example Show that $f(x) = 2x^5 - x^4 + 2x^3 - 2x^2 + 4x - 4$ has no real zero greater than 2

Lesson: Synthetic Division and the Remainder Theorem

The outcome of division can be restated using the division algorithm.

State the division algorithm:

Show how we use polynomial long division with $(x^2 - 3x - 1) \div (x + 2)$, and then restate the result using the format of the division algorithm.

Now show how to do the operation $(x^2 - 3x - 1) \div (x + 2)$ with synthetic division:

Restate this result using the division algorithm:

Example Use synthetic division to perform: $\frac{4x^3 - 3x^2 - 7}{x + 1}$. Rewrite the result using the division algorithm.

So what are 3 keys about using synthetic division to remember?

What is the remainder theorem?

Example Use the remainder theorem to find the desired functional values: $f(x) = -3x^4 + 5x^3 - 30x + 20$, $f(2)$. What is the equivalent division problem associated with this? Write the result of your division using the division algorithm.

Example Use the remainder theorem to find the desired functional values: $f(x) = 3x^2 + 5, k = 3 - i$

Lesson: Synthetic Division and Potential Zeros of Polynomials

What is a “zero” of a polynomial?

How does the remainder theorem help us to determine if we have a zero?

What are some other terms that we use for “zero” of a polynomial?

Example Determine if the given k is a zero of the polynomial. If it is not, then what is the value of $f(k)$?

a. $f(x) = x^2 - 4x - 5; k = 5$

b. $f(x) = 2x^3 - 3x^2 + 5; k = 1$

c. $f(x) = x^2 - 2x + 2; k = 1 + i$

Lesson: The Factor Theorem

What does the factor theorem say?

Example Determine whether $(x - 1)$ is a factor of each polynomial:

a. $f(x) = 9x^4 - 13x^2 + 4$

b. $f(x) = 4x^3 + 10x^2 + 2x - 4$

Given the above, what can we say is also a zero for example a?

Example Factor $f(x) = 3x^3 + 4x^2 - 17x - 6$ into linear factors given that -3 is a zero.

Example Factor $f(x) = 2x^3 - x^2 - 8x + 4$ into linear factors given that $\frac{1}{2}$ is a zero.

Lesson: The Rational Zeros Theorem

The rational zeros theorem is a way to determine *possible* candidates for *rational* zeros of a polynomial function.

State the Rational Zeros Theorem:

Example Find all the possible rational zeros for $f(x) = 2x^3 - x^2 - 8x + 4$ and then factor into linear factors.

Example Find all the possible rational zeros for $f(x) = 5x^4 - 19x + x^2 + 31x + 6$ and then factor into linear factors.

Lesson: The Fundamental Theorem of Algebra and the Number of Zeros Theorem

What does the Fundamental Theorem of Algebra state?

What is the number of zeros theorem?

Example What is the maximum number of zeros that each polynomial could possibly have?

a. $f(x) = x^2 - 3x + 1$

b. $f(x) = x^{17} - 3x + 1$

c. $f(x) = 9$

Example Find a polynomial $f(x)$ that has zeros $-3, 3$, and 1 , where $f(2) = -10$

Example Find a polynomial $f(x)$ that has zeros $6, -3$, and -2 , where $f(1) = 20$

Lesson: Conjugate Zeros Theorem

Before we can discuss the theorem, let's review a few properties of conjugates.

What is the conjugate of the following:

a. $3 + 2i$

b. $5 - 3i$

c. $-7 + i$

What is another way that we denote conjugates? Summarize the explanation in the video. Give an example of how this notation works.

What are some properties of conjugates?

What is the conjugate zeroes theorem?

Example Find a polynomial function of least degree with only real coefficients and zeros 2 and $(3 - i)$.

Example Suppose $k = -i$ is one zero of $f(x) = x^4 + 26x^2 + 25$. What are the other zeros?

Example Factor $f(x) = 2x^3 + (3 - 2i)x^2 + (-8 - 5i)x + (3 + 3i)$ into linear factors given $k = 1 + i$ is a factor.

Before you begin, what is unique about this problem?

Lesson: Finding Zeros of a Polynomial Function

In previous videos we discussed:

- The factor theorem
- The rational zeros theorem
- The fundamental theorem of algebra
- The number of zeros theorem
- The conjugate zeros theorem

Using all of these theorems, we can combine them to find the zeros of *any* polynomial.

What can we say about a polynomial function of odd degree with real coefficients?

What can we say about a polynomial function of even degree with real coefficients?

What if a polynomial doesn't have only real coefficients?

Example Find all zeroes of the following polynomials.

a. $f(x) = x^3 - 21x - 20$

b. $f(x) = 2x^4 - x^3 + 7x^2 - 4x - 4$

Examples: Finding Zeros of a Polynomial Function

Find all the zeros of the polynomials.

a. $f(x) = x^4 + 13x^2 + 36$

b. $f(x) = 2x^4 - 11x^3 + 4x^2 + 29x + 12$

c. $f(x) = x^4 - 7x^3 + 17x^2 - x - 26$

Lesson: Descartes' Rule of Signs

Let $f(x)$ be a polynomial function with real coefficients and a nonzero constant term, with terms in descending powers of x . Explain how to use Descartes' Rule of Signs:

Example Determine the number of different possibilities for the number of positive, negative, and nonreal complex zeroes:

a. $f(x) = 3x^5 + 4x^4 - x^3 + 2x - 1$

b. $f(x) = x^6 + x^5 - x^4 + x^3 + x^2 - x + 1$

c. $f(x) = 3x^5 - 7x^3 + 6x + 8$

Lesson: The Basic Graphs of Rational Functions


A rational function is a function of the form:

$$\frac{f(x)}{g(x)}, g(x) \neq 0$$

Where $f(x)$ and $g(x)$ are polynomials.

There are 2 graphs that we view as the most basic version of rational functions.

Make a table and graph $f(x) = \frac{1}{x}$




What is the domain and range?

Where does the graph not exist (ie where is it discontinuous)

What are the y-axis and x-axis for this graph?

What kind of symmetry does this graph have?

Make a table and graph $f(x) = \frac{1}{x^2}$



What is the domain and range?


Where does the graph not exist (ie where is it discontinuous)

What are the y-axis and x-axis for this graph?


What kind of symmetry does this graph have?

Example Describe the translations for the following graphs, and then draw an approximate sketch of the graph.

a. $f(x) = \frac{1}{(x+1)^2} - 2$



b. $g(x) = -\frac{2}{x} + 1$



Sketch what happens for odd powers of n for: $f(x) = \frac{1}{x^n}$, ie just draw a sketch of $y = \frac{1}{x^3}$, $y = \frac{1}{x^5}$, $y = \frac{1}{x^7}$

Sketch what happens for higher even powers of n for: $f(x) = \frac{1}{x^n}$, ie just draw a sketch of $y = \frac{1}{x^4}$, $y = \frac{1}{x^6}$, $y = \frac{1}{x^8}$

Lesson: Asymptotes of Rational Functions

What is an informal definition of asymptotes?

Is it possible that a graph would graph an asymptote?

Here are the formal definitions of asymptotes:

A graph $f(x) = \frac{p(x)}{q(x)}$ ($q(x) \neq 0$) has a

- **Vertical asymptote** if $|f(x)| \rightarrow \infty$ as $x \rightarrow a$. The asymptote is $x=a$.
- **Horizontal asymptote** if $f(x) \rightarrow b$ as $|x| \rightarrow \infty$. The asymptote is $y=b$.

What is super important about asymptote?

Where do you typically find vertical asymptotes?

How can you determine horizontal asymptotes? Describe the two ways, and also explain how oblique asymptotes. (In other words, explain what we talked about in the video).

Examples Find any vertical, horizontal, or oblique asymptotes.

a. $f(x) = \frac{x+2}{x^2+7x+12}$

b. $f(x) = \frac{3x+1}{4x-2}$

c. $f(x) = \frac{x^2+3x+2}{x^2-9}$

d. $f(x) = \frac{x^2+x}{x+3}$

Examples: Finding Oblique Asymptotes

Find all the asymptotes for the following functions.

a. $f(x) = \frac{x^2+2x}{2x+1}$

b. $f(x) = \frac{x^2+2}{x+1}$

Lesson: Graphing Rational Functions

To learn how to graph rational functions, we will outline the steps and then graph. **Clearly write directions for each step as we go through this process!!!**

Graph $f(x) = \frac{x+3}{x^2+7x+10}$. Outline all the relevant steps discussed in the video.

Graph $f(x) = \frac{x+1}{x-3}$. Outline all the relevant steps discussed in the video.

Graph $f(x) = \frac{x^2+1}{x-3}$. Outline all the relevant steps discussed in the video.

Examples: Graphing Rational Functions

Graph $f(x) = \frac{(x+2)^2}{(x-1)(x+3)}$. Outline all the relevant steps discussed in the video.

Graph $f(x) = \frac{x}{x^2-4}$. Outline all the relevant steps discussed in the video.

Graph $f(x) = \frac{2x^2-3x-1}{x-2}$. Outline all the relevant steps discussed in the video.

Examples: Equations of Rational Functions

Copy the graphs from the video and then describe how to find the equations of those graphs.

Variation

Video Notes

The term “variation” refers to a particular way of describing how certain things relate to one another.

There are three types of variation we will look at. Write down the definitions for each, you will remember it better!

1. Direct Variation:

2. Inverse Variation

3. Joint Variation

It’s also possible to combine different types of variation.

The first thing we have to do is learn how to interpret and write variation equations.

Examples Write the variation equation:

- a. P varies directly with Q

- b. J varies inversely with L

- c. H varies jointly with A and B

- d. Z varies inversely with the square of R

- e. M varies directly as the square root of N

- f. Y varies jointly with the cube of M and the square root of L

- g. M varies directly as L and inversely as N

Now we can take a look at how to solve general variation equations.

Example Write down the detailed steps / directions for this example.

J varies directly with L. If $J = 12$ when $L = 4$, find J when $L = 7$.

Step 1:

Step 2:

Step 3:

Step 4:

Step 5:

Examples Using the steps outlined from the previous example, try the following:

- a. M varies inversely as the square of L . If $M = 12$ when $L = 2$, find M when $L = 4$.
- b. A varies jointly with B and C . If $A = 30$ when $B = 2$ and $C = 5$, find A when $B = 3$ and $C = 4$.
- c. L varies directly as M and inversely as N . If $L = 15$ when $M = 5$ and $N = 2$, find L when $M = 7$ and $N = 3$.

Example Jada's weekly income varies directly with the amount of hours that she works. Last week, she earned \$570 for working 38 hours. This week she worked 36 hours, how much will she earn?

Lesson: A Review of Variation

Describe the 3 types of variation and outline any patterns that might help you remember.

Direct Variation

Inverse Variation

Joint Variation

Combined Variation

Example If y varies directly with x , and $y=15$ when $x=3$, find y when $x=6$

Example If a varies jointly with b and c , and $a=40$ when $b=2$ and $c=5$, find a when $b=3$ and $c=2$.

Example If p varies inversely with m and directly with the square of t , and $p=32$ when $t=2$ and $m=4$, find p when $t=3$ and $m=9$

Example Rewrite the following situations as variation.

a. $C = 2\pi r$, when C is the circumference of a circle and r is the radius.

b. $r = \frac{d}{t}$, where r is the speed when traveling d miles over t hours.

c. $f = \frac{mv^2}{r}$, where f is the centripetal force of an object of mass m moving along a circle with radius r at velocity v.

Examples: Applications of Variation

The circumference of a circle varies directly as the radius. A circle of radius 8in has a circumference of 50.27in. Find the circumference of the circle if the radius is 5in.

The current in a simple electrical circuit varies inversely as the resistance. If the current is 30 amps when the resistance is 10ohms, find the current if the resistance is 5ohms.


The volume of a gas varies inversely as the pressure and directly as the temperature. If a gas occupies a 2.6L volume at 600 kelvins (kelvin is a temperature unit) and a pressure of 24 newtons, find the volume at 240 kelvins and a pressure of 32 newtons.

Lesson: Exponential Functions


An exponential function is a function of the form _____ where $a > 0, a \neq 1$.

Examples Graph the following functions, then determine their domain and range.

a. $f(x) = 2^x$




b. $g(x) = \left(\frac{1}{2}\right)^x$




Using the graphs above, summarize some of the characteristics of these graphs:

The number e is an irrational number that is frequently used in math. What is an approximation of e ?

Example Graph $f(x) = 2^x$, $g(x) = e^x$, $h(x) = 3^x$ all on the same graph below for comparison:



Translations work similarly to other graphs we have discussed. Using the graph of $f(x) = 2^x$ as a “base graph”, draw the following graphs together on the same graph: $f(x) = 2^x$, $g(x) = 2^x + 3$, $h(x) = 2^{x-1}$



Lesson: Solving Exponential Equations

In this lesson, we will solve equations that look like this:

$$3^x = 27 \quad \text{or} \quad 4^{x+3} = 8^{x-2}$$

To do this, we will write directions around the following example: $2^{3x} = 8^{x+1}$

Examples Solve:

a. $49^{x-3} = 7^{2x}$

b. $9^{4x} = 27^{x+4}$

Examples Solve:

a. $5^{x-2} = \frac{1}{25}$

b. $\left(\frac{3}{5}\right)^{2x} = \left(\frac{27}{125}\right)^{x+2}$

c. $27^{3x+1} = 9^{4x+2}$

Lesson: A Review of Exponential Functions and their Graphs

What is the form of a general exponential equation?

Example Let $f(x) = 3^x$ and $g(x) = \left(\frac{1}{2}\right)^x$. Find the following.


a. $f(-2)$

b. $f\left(\frac{3}{2}\right)$

c. $g(-3)$

d. $g(3.2)$

Sketch the graph of $f(x) = 2^x$ and create a table to go with it.




When will an exponential function $f(x) = a^x$ take this shape?

What is the x-axis for this graph?

What are 3 standard points to note here?

What is the domain and range?

Sketch the graph of $f(x) = \left(\frac{1}{2}\right)^x$ and create a table to go with it.



When will an exponential function $f(x) = a^x$ take this shape?


What is the x-axis for this graph?

What are 3 standard points to note here?


What is the domain and range?

Examples Describe the translations and sketch the following graphs.

a. $f(x) = -3^x + 1$



b. $f(x) = \left(\frac{1}{2}\right)^{x+1} - 2$



Lesson: Solving Basic Exponential Equations

Outline the general steps to solve $3^{x+1} = 81$.

Outline the general steps to solve $4^{x-2} = 8^{x+1}$.

Examples Solve the following.

a. $27^{2x} = 9^{x-1}$

b. $e^{3-x} = (e^2)^{-x}$

c. $(\sqrt{3})^{x+3} = 9^x$

Examples Solve the following.

a. $x^{\frac{3}{2}} = 8$

b. $x^{\frac{5}{2}} = 4$

c. $x^{-4} = \frac{1}{16}$

Lesson: Compound Interest

What is the formula for compound interest and what does it mean?

Example Suppose \$2000 is invested into an account paying 2.5% interest per year compounded monthly. How much is in the account after 10 years, and how much interest is earned in that time?

What else do we call the A from the compound interest formula?

What else do we call the P from the compound interest formula?

Example In 5 years, you want to have \$8000 in cash to buy a car (including taxes). Let's use savings to help us reach this goal.

a. You find an account with a 3.25% interest rate compounded biannually (2x a year). What amount would you have to deposit today to ensure you have \$8000 in 5 years?

b. Let's pretend that as of today, you have \$5000. What interested rate would you need to ensure that this will grow to \$8000 in 5 years?

What is the number e ?


What is the formula for continuous compounding?

Example Suppose \$2000 is invested into an account paying 2.5% interest per year compounded continuously. How much is in the account after 10 years, and how much interest is earned in that time?

Lesson: Intro to Logarithmic Functions

First, we want to review how to function inverses of functions. Find the inverse of: $f(x) = 2x + 8$

Graph the exponential function $f(x) = 3^x$ below:



Recall the exponential functions are one-to-one, which means that they have inverses. Attempt to find the inverse of $f(x) = 3^x$ below:

As we can see, there is an issue with finding the inverse – we have no way to solve for y ! As such, we need new mathematics to help solve this.

Write down the definition of a logarithm:

Examples Write the following in exponential form:

a. $\log_3 81 = 4$

c. $\log_5 125 = 3$

b. $\log_2 \frac{1}{8} = -3$

d. $\log_9 9 = 1$

Examples Write the following in logarithmic form:

a. $2^4 = 16$

c. $6^{-2} = \frac{1}{36}$

b. $3^0 = 1$

d. $8^{\frac{1}{3}} = 2$

Now we can finally determine what is the inverse of $f(x) = 3^x$:

There are a few common logarithms:

- $\log x$ means:

- $\ln x$ means:

There are also 2 key properties of logarithms:

1. $\log_a a = 1$: explain why this makes sense

2. $\log_a 1 = 0$: explain why this makes sense:

Lesson: Basics of Solving Logarithmic Equations and Evaluating Logarithms

We want to solve equations that look like:

$$\log_3 x = 4 \quad \text{or} \quad \log_x 36 = 2$$

Before you begin, you should already be able to do the following:

a. Put 3^5 in logarithmic form:

b. Put $\log_5 5 = 1$ in exponential form:

Now we can explain how to solve logarithmic equations using the example $\log_3(2x + 1) = 3$

Example Solve: $\log_x 16 = 4$

Example Solve:

a. $\log_9 x = 2$

b. $\log_6(2x + 4) = 2$

Now we can also discuss how to evaluate logarithms. Explain how to evaluate: $\log_3 27$:

Examples Evaluate:

a. $\log_5 \frac{1}{25}$

b. $\log 100$

Examples Evaluate:

a. $\ln e$

c. $\log_{25} 5$

b. $\log_3 27$

d. $\log \frac{1}{10}$

Lesson: Review of the Basics of Logarithmic Functions

Explain how to find the inverse of $f(x) = 3^x$, and explain how this relates to logarithms.

Define a logarithm:

Example Write the following in logarithmic form.

a. $3^2 = 9$

b. $5^0 = 1$

c. $4^{-2} = \frac{1}{16}$

Example Write the following in exponential form.

a. $\log_3 1 = 0$

b. $\log_5 125 = 3$

c. $\log_3 81 = 4$


Example Show how to solve the following basic logarithmic equations.

a. $\log_x 8 = 3$

b. $\log_3 x = -2$

c. $\log_{49} 7 = x$

Create a table and show how to graph $f(x) = \log_2 x$




What 3 points can you easily find on the graph?

When does the graph take a shape like this for $f(x) = \log_a x$?

What is the y-axis for this graph?

What are the domain and range?

Create a table and show how to graph $f(x) = \log_{\frac{1}{2}} x$




What 3 points can you easily find on the graph?

When does the graph take a shape like this for $f(x) = \log_a x$?

What is the y-axis for this graph?

What are the domain and range?

Examples Graph the following: $g(x) = \log_2(x - 1) + 1$



Lesson: How to Graph a Logarithm


Before we begin, you should be able to do the following:

a. Convert to logarithmic form: $25 = 5^2$

b. Convert to exponential form: $\log_2 8 = 3$


c. Find the inverse of $y = 2^x$

Now we can explain how to graph $y = \log_3 x$



For comparison, also graph $f(x) = 3^x$ on the same graph above.

Example Graph $y = \log_{\frac{1}{2}} x$ below:



Lesson: Review of the Basics of Properties of Logarithms

List the major properties of logarithms.

Explain what the common logarithms mean:

$\log x =$

$\ln x =$

Examples Expand the following as much as possible.

a. $\log_3 \left(\frac{3x^3y^7}{\sqrt{z}} \right)$

b. $\ln \left(\sqrt[3]{x^2y} \right)$

c. $\log_3 \sqrt{6}$

d. $\log_5 \frac{25x^3y^2}{p^7r^3}$

Example Write as a single logarithm.

a. $\log(x + 2) + \log(x - 2) - \log(x + 3)$

b. $3 \log_2 3 + 2 \log_2 y - 5 \log_2 x$

c. $\frac{1}{2} \log_5 16 + 3 \log_5 y^2 - \log_5 x^7 - \log_5 y$

d. $4 \ln 2 + \ln x^6 - 2 \ln y^3 - 3 \ln x^2$

Lesson: Change of Base Theorem

State the change of base theorem:

Example Show how to evaluate the following:

a. $\log_5 17$

b. $\log_2 3$

c. $\log_{\frac{1}{2}} 5$

Lesson: Solving Exponential Equations

Outline the steps to solve: $2^{x+3} = 7^{3x}$

Examples Solve the following.

a. $e^{x^2} = 100$

b. $e^{2x-3} * e^{3x} = 5e$

Examples Solve.

a. $3(2)^{x+2} + 6 = 105$

b. $e^{2x} + 8e^x + 15 = 0$

c. $3^{2x} + 2(3^x) = 24$

Lesson: Solving Logarithmic Equations

Outline the steps to solve: $\log(3 - x) = 0.75$

Compare these steps to: $\log(x + 20) = \log(x + 10) + \log(6)$. How are these problems different?

Examples Solve the following.

a. $\log_4(x^3 + 37) = 3$

b. $\log x + \log x^3 = 4$

Examples Solve the following.

a. $\log(x^2 - 4) - \log(x - 2) = \log(8)$

b. $\log(5 + 4x) - \log(3 + x) = \log 3$

Lesson: Solving Exponential and Logarithmic Formulas

Examples Solve for the desired variable.

a. for b: $p = a + \frac{c}{\ln b}$

b. for n: $A = \frac{Pr}{1-(1+r)^{-n}}$

c. for A: $\log A = \log B - C \log D$

d. for x: $D = 100 + 5 \log x$

Lesson: Models of Exponential Growth and Decay

Explain the equation for the general model:

Some examples of growth can come from populations or investments. Some examples of decay can come from half-life/radioactive decay, or carbon dating.

Define the following:

Doubling time:

Half-life:

What is the formula for continuous compounding?

Example A sample of 400g of radioactive lead decays to polonium according to the function: $A(t) = 400e^{-0.03t}$.

a. Find the amount of lead remaining after 5 years.

b. Find the half-life.

Example The amount of carbon-14 present in a paint after t years is given by $y = y_0e^{-0.0002t}$. The paint contains 37% of its carbon-14. How old are the paintings?

Example Find the doubling time of an investment earning 5% interest if interest is compounded continuously.

Example During an epidemic, the number of people who have never had a particular virus and are not immune decreases exponential, where this decline is modeled by the function: $f(x) = 20,000e^{-0.025t}$, where t is time in days.

a. How many people were susceptible at the beginning of the epidemic?

b. How many people are susceptible after 2 weeks?

Lesson: A Review of Angles and Degrees

Draw, label, and define how to make an angle:

Show how to create an angle of positive versus negative measure:

Degree is generally how we measure angles.

Draw the complete rotation of a ray. How many degrees is this?

Define:

- Acute angle
- Right angle
- Obtuse angle
- Straight angle

Define:

- Complementary angles
- Supplementary angles

Example Find the complement and supplement of a 55° angle.

Example Find the measures of the marked angles shown in the video.

Examples: Converting Between Minutes and Seconds

Examples Convert the following to degrees.

a. $23^{\circ}15'32''$

b. $38^{\circ}27'15''$

Examples Convert the following to degrees, minutes, and seconds.

a. 32.713°

b. 72.17°

Examples: Adding and Subtracting Degrees, Minutes, and Seconds

Add or subtract.

a. $45^{\circ}15' + 23^{\circ}57'$

b. $23^{\circ}18'29'' + 14^{\circ}19'47''$

c. $94^{\circ}16' - 27^{\circ}29'$

d. $72^{\circ}18'15'' - 15^{\circ}20'30''$

Examples: Applications of Minutes and Seconds

Example A turntable makes 45 revolutions in a minute. How many revolutions is that per second?

Example A windmill makes 80 revolutions per minute. How many revolutions is that per second?


Example Wind power is caused by a turbine spinning 10 to 20 revolutions per minute. How many revolutions does it make per second?

Lesson: Angle Measure with a Clock


Explain the basics of angle measure, ie where is 90° , 180° ?

Each number represents how many degrees? SHOW how to actually determine this?


Example Find in the clock below, and show how to find the angle measure for 5:00. Draw pictures to help illustrate how this works, pictures are key to remembering!!!



Example Find in the clock below, and show how to find the angle measure for 8:20. Draw pictures to help illustrate how this works, pictures are key to remembering!!!




Example Find in the clock below, and show how to find the angle measure for 10:05. Draw pictures to help illustrate how this works, pictures are key to remembering!!!




Examples: Angle Measure from a Clock

Find the angles associated with each time.


a. 6:25



b. 3:15



c. 2:38



Lesson: Standard Position and Coterminal Angles

When is an angle in standard position? Draw an example.

Show some examples of how we can determine angle measure by quadrant.

What are quadrantal angles? Draw a picture to illustrate.

What makes two angles coterminal? Draw a picture to explain.

Example Find the angle of least positive measure that is coterminal with each angle:

a. 715°

b. -80°

Example How can you find all angles coterminal with 45° ? Prove it by filling in the following table.

Value of n	Angle coterminal with 45°
2	
0	
-1	

Examples Fill in the table below for which angles are coterminal with the quadrantal angles

Quadrantal Angle	Coterminal Angles
0°	
90°	
180°	
270°	

Lesson: Radian Measure

Radian measure is another way to describe angle measure, as opposed to degrees.

What does radian measure allow us to do?

What is the definition of a radian?

Are radians a unit of measure like feet or meters? Circle:

Yes

No

What is the formula for the distance around a circle?

What does this formula show?

Write a way to relate radians to degrees:

So what is π radians?

Show a conversion rate between radians and degrees:

Examples Convert each degree to radians:

a. 35°

b. -175°

c. 27.8°

Convert each radian measure to degrees:


a. $\frac{7}{4}\pi$

b. $-\frac{2\pi}{3}$

c. 7

If no unit of angle measure is specified, what should you assume?

Below we will construct an illustration of all the common degree measures we have discussed, and then write their radian equivalent. Take your time to copy this perfectly!



You will need to know all of those common degrees AND radian measures for the exam!

You may want to reference cheat sheets to help you with the next few problems.

Examples Find the value of each function.

a. $\sin \frac{5\pi}{6}$

b. $\tan \frac{3\pi}{4}$

c. $\csc \frac{5\pi}{3}$

d. $\cos \left(-\frac{2\pi}{3} \right)$

Lesson: Arc Length on a Circle and Some Applications

We can determine the formula for the arc length of a circle from the definition of radians.

Regarding radians, what does θ equal? Show how this relates to arc length.

What is key about the arc length formula?

Examples A circle has radius 22.30 cm. Find the length of the arc intercepted by a central angle having each of the following measures.

a. $\frac{5\pi}{6}$ radians

b. 66°

Examples A circle has radius 15.22 cm. Find the length of the arc intercepted by a central angle having each of the following measures.

a. $\frac{3\pi}{8}$

b. 125°

What is the definition of latitude?

Example The latitude of Panama City, Panama is 9° N and the latitude of Pittsburgh, PA is 40° N. Find the north-south distance of these 2 cities. The radius of the Earth is 6400 km.

Example A rope is being wound around a drum with a radius of 0.7525 ft. How much rope will be wound around the drum if the drum is rotated through an angle of 42.25° ?

Example Two gears are adjusted so that the smaller gear drives the larger one. If the smaller gear rotates through an angle of 150° , through how many degrees will the larger gear rotate? Refer to the photo in the video.

Examples: Applications using Arc Length

Charleston, SC and Toronto, Canada, are 1100 km apart and lie on the same north-south line. If the latitude of Charleston is 33° N, what is the latitude of Toronto?

Find the radius of the larger wheel in the figure if the small wheel rotates 75.0° when the large wheel rotates 45° .

How many inches will a small weight rise if a pulley is rotated through an angle of $65^{\circ}15'$? Through what angle, to the nearest minute, must the pulley be rotated to raise the weight 4 inches?

Lesson: Area of a Sector of a Circle and Some Applications

What do we define as a sector of a circle?

What is the formula for the area of a section of a circle with radius r ?

Examples Find the area of a sector of a circle having the given radius and central angle θ . Answers should be to the nearest tenth.

a. $r = 11.5 \text{ ft}, \theta = \frac{5\pi}{6} \text{ radians}$

b. $r = 13.5 \text{ m}, \theta = 125^\circ$

Example Find the measure (in radians) of a central angle of a sector of a circle of area 25 in^2 in a circle of radius 2.5 in.

Example A center-pivot irrigation system provides water to a sector-shaped field as shown in the figure in the video. Find the area of the field if $\theta = 35.0^\circ$ and $r = 155 \text{ yd}$.

Example A circular sector has an area of 60 in^2 . The radius of the circle is 6 inches. What is the arc length of the sector?

Lesson: Linear Speed and Angular Speed and Applications

What is the formula for linear speed?

When are some situations where you might use linear speed?

What is the formula for angular speed?

When are some situations where you might use angular speed?

Examples Use the formula $\omega = \frac{\theta}{t}$ to find the value of the missing variable.

a. $\omega = \frac{\pi}{3}$ radians per second, $t = 6$ seconds

b. $\theta = 3.721$ radians, $t = 18.34$ seconds

Examples Use the formula $v = r\omega$ to find the value of the missing variable.

a. $r = 10$ m, $\omega = \frac{9\pi}{5}$ radians per second

b. $v = 10$ m per second, $\omega = \frac{3\pi}{2}$ radians per second

Example Suppose a point P is on a circle with a radius of 5 cm, and the ray OP is rotating with an angular speed of $\frac{\pi}{10}$ radian per second.

a. Find the angle generated by P in 3 seconds.

b. Find the distance traveled by P along the circle in 3 seconds.

c. Find the linear speed of P in cm per second.

The formula $\omega = \frac{\theta}{t}$ can be rewritten as $\theta = \omega t$. We can then use the formula $s = r\theta$ to create a different formula. Show the new formula we can create:

Examples Use the formula above to find the value of the missing variable.

a. $r = 5 \text{ cm}$, $\omega = \frac{2\pi}{3}$ radians per second, $t = 8$ seconds

b. $s = \frac{3\pi}{2} \text{ km}$, $r = 4 \text{ km}$, $t = 3$ seconds

Example The orbit of the earth around the sun is almost circular. Suppose that the orbit has a radius of 93 million miles.

a. We know that a year is 365 days. Find the angle formed by the Earth's movement in one day.

b. Give the angular speed in radians per hour.

c. Find the linear speed of the Earth in miles per hour.

Examples: Applications of Linear Speed and Angular Speed

Example Find the angular speed of the minute hand of a clock.

Example Find the linear speed of the tip of the second hand of a clock, if the hand is 25 mm long.

Example A bicycle has tires with a 12-inch radius. The tires are turning at the rate of 200 revolutions per minute. How fast is the bike traveling in miles per hour?

Example A pulley has a radius of 13.23 cm. It took 15 seconds for 45 cm of belt to go around the pulley.

a. Find the linear speed of the belt in cm per sec.

b. Find the angular speed of the pulley in radians per second.

Example A thread is being pulled off a spool at a rate of 55.4 cm per second. Find the radius of the spool if it makes 155 revolutions per minute.

Lesson: Angle Relationships

Draw and explain what vertical angles are:

Draw and explain what makes a transversal line. Draw a big picture, you will use it for the next part of your notes:

Label the diagram and note the relationships between the different angles below:

Example Draw and label the lines from the video. Find the measures of angles 1, 2, 3, and 4.

Example Draw and label the lines from the video. Find the measures of angles 1, 2, 3, and 4.

Lesson: Review of Triangles and Similar Triangles

The sum of angle measures of any triangle is:

Draw the different types of triangles:

Examples Draw the triangles in the example, then classify the triangles as acute, right, or obtuse. Also classify as equilateral, isosceles, or scalene.

What makes 2 triangles similar?

What makes 2 triangles congruent?

What is the key relationship between similar and congruent triangles?

What are the 2 conditions for similar triangles?

In other words, triangle ABC is similar to triangle DEF if:

Example Draw the similar triangles in the video, and then find any missing angle measures.


Example Draw the similar triangles in the video, and then find any missing side lengths.

Example Draw the similar triangles in the video, and then find any missing measures.


Examples: Similar Triangles

For each example, assume the triangles are similar. Triangles are not drawn to scale. Find any missing angle or side measures.


a.



b.



c.



Lesson: Right-Triangle Based Definitions of Trigonometric Functions and Cofunctions

Instead of using standard position to describe trig functions, we can alternatively use right triangles.

Draw a right triangle to orient yourself.

What does A represent?

Show how to define the 6 trig functions with this right triangle.

1.

2.

3.

4.

5.

6.

What mnemonic device can help you remember the trig functions?

Example Draw the triangle in the video, and then show how to find the values of sine, cosine, and tangent.

Example Using the same example above, show how to find the values of cosecant, secant, and cotangent.

Draw a new right triangle, and show an alternative angle measure we can find from this picture:

These relationships are called the **cofunction identities**, summarize these below:

Examples Write each function in terms of its cofunction

a. $\sin 10^\circ$

b. $\cot 65^\circ$


c. $\csc 34^\circ$

Example Find one solution for each equation. Assume all angles involved are acute angles.

a. $\cos \theta = \sin (2\theta - 30^\circ)$

b. $\sec(\theta + 15^\circ) = \csc(2\theta + 25^\circ)$

Lesson: More on Trig Function Values as Angles Change and Special Angles



Explain how angle A changes in these pictures:

Examples Determine whether each statement is true or false.

a. $\tan 25^\circ < \tan 22^\circ$

b. $\cos 25^\circ < \sin 25^\circ$

There are a few angles that occur frequently in trig and have some *nice* properties that we like to call them out. Draw out a triangle where each angle is 60° and describe any other properties:

If you bisect this triangle, what happens? Draw a picture to illustrate, and then describe the trig functions associated with it:

Finally, draw an isosceles right triangle and describe any special characteristics with it:

We have now examined some *extremely* common trig measures that will come up all the time in trig. They are so important that we will create a separate reference sheet for them.

Read this! I highly recommend that you remove the last page from this packet, and instead put it at the very front of your notes so that you can easily reference it. If you are moving on to calculus, KEEP THIS PAGE IN GOOD CONDITION FOR THE NEXT FEW YEARS. You will thank yourself later that you have this reference as these angles will come up over and over again, and students frequently forget them. I will occasionally create these handouts and make these suggestions to you as we go along in trig – you will thank yourself (and me) later if you keep these. It is extremely common for students to forget some of KEY parts of trig and then struggle in calculus – I don't want this to happen to you!

Some Function Values of Special (Common) Angles

θ	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\cot \theta$	$\sec \theta$	$\csc \theta$
0°						
30°						
45°						
60°						
90°						

Lesson: The Unit Circle and Circular Functions

Explain how we create the unit circle:

What is the equation for the unit circle?

Recall the formula for arc length is $s = r\theta$. When the radius of the circle is 1, what does this formula become?

Why does this matter / what is cool about this?

Since $r=1$ on the unit circle, what can we say about our previously studied trig functions? Just summarize what we discussed in the video as best as you can.

Write out the **circular functions**

1.

2.

3.

4.

5.


6.

Explain the symmetry of the unit circle:

If a point (a, b) is on the unit circle, what other points are guaranteed to be on the unit circle?

How can we use symmetry with drawing the unit circle? How is it handy?

Label the unit circle. Note: you will have to have this memorized! These are the standard angle measures that you'd be expected to do without a calculator!



Now, going back to the circular functions we discussed, how do you view the x-coordinates?

How do you view the y-coordinates?

Examples Use the unit circle to find the values of the following:

a. $\sin \frac{\pi}{3}$

b. $\cos \frac{5\pi}{4}$

c. $\sin \frac{11\pi}{6}$

Lesson: Arc Length on a Circle and Some Applications

Using the unit circle, we can also determine the domains of our circular functions.

What is the domain of sine and cosine, and why?

What is the domain of tangent and secant, and why?

What is the domain of cotangent and cosecant, and why?

Lesson: Values of Circular Functions

We have previously discussed the unit circle (you will want to have it handy). One the most important things about the unit circle is that it will quickly tell you the value of the sine and cosine functions.

On the unit circle, what does the y-coordinate represent?

On the unit circle, what does the x-coordinate represent?

When we think about the values of these functions, what do we assume?

What does this mean for your calculator?

Examples Use the unit circle to find the values of:

a. $\sin \frac{\pi}{2}$

b. $\cos \frac{\pi}{2}$

c. $\tan \frac{\pi}{2}$

Examples Use the unit circle to find the values of:

a. $\sin \frac{5\pi}{4}$

b. $\cos \frac{5\pi}{4}$

c. $\tan \frac{5\pi}{4}$

Example Compare methods. Use reference angles and degrees to find the exact value of $\sin \frac{2\pi}{3}$, then confirm your answer with the unit circle.

For approximate values, we still must use a calculator. Remember, when you are not specifically told you are using degrees, then you assume you are using radians.

What must you ensure with your calculator?

Examples Use a calculator to approximate the function values.

a. $\cos(1.72)$

b. $\sin(-3.54)$

c. $\cot(1.438)$

Lesson: Values of Inverse Circular Functions

What do the functions $\sin^{-1} x$, $\cos^{-1} x$, $\cot^{-1} x$, etc. represent?

Your calculator makes a few key assumptions if it uses these functions (the reasons why are explained later):

1. For all x in $[-1, 1]$, using $\sin^{-1} x$ with a calculator will return a value between:

2. For all x in $[-1, 1]$, using $\cos^{-1} x$ with a calculator will return a value between:

3. For all real numbers x , using $\tan^{-1} x$ with a calculator will return a value between:

Sometimes you will be asked to find answers in different quadrants than what your calculator will return!

Example Solve for s , given that s is in the interval $\left[0, \frac{\pi}{2}\right]$: $\sin s = 0.3311$

Example Solve for s , given that s is in the interval $\left[0, \frac{\pi}{2}\right]$: $\tan s = 0.2155$

Example Solve for s , given that s is in the interval $\left[\pi, \frac{3\pi}{2}\right]$: $\tan s = \sqrt{3}$

Example Solve for s , given that s is in the interval $[0, 2\pi)$: $\cos^2 s = \frac{1}{2}$

Lesson: The Unit Circle and Circular Functions as Line Segments

The unit circle is really just an extension of how we have previously thought about trig functions as line segments / right triangles.

Draw the illustration in the video, then show how these sides relate to our trig functions.

From your illustration, how do we define the following?

$\cos \theta =$

$\tan \theta =$

$\csc \theta =$

$\sin \theta =$

$\cot \theta =$

$\sec \theta =$

Example Using your illustration from the previous page, suppose the measurement of $\angle TVU$ is 45° . Find the lengths of the following line segments:

OQ

OU

OV

VR

US

PQ

Lesson: The Pythagorean Theorem, The Distance Formula, and Intro to Trig Functions

State the Pythagorean theorem:

State the distance formula:

How does the Pythagorean theorem relate to the distance formula?

Now we will introduce **trigonometric functions!** Explain what makes trig functions unique:

How do trig functions relate to the distance formula? Draw the picture associated with this.

Explain the 6 trig functions and how they relate to what we discussed above.

Sometimes you will see expressions such as $\sin^2 90^\circ$. What does this mean?

Example The terminal side of an angle θ in standard position passed through the point $(-12, -5)$. Find the values of the 6 trig functions of angle θ .

Example Find the 6 trig function values of an angle θ in standard position if the terminal side of θ is defined by

$$3x + 5y = 0, x \geq 0$$

Example Suppose the point (x, y) is in the indicated quadrant. Determine whether given ratio is positive or negative.

$$III, \frac{y}{r}$$

Examples: Finding Values of Trig Functions

For each given point, sketch the angle of θ in standard position such that θ has the least positive measure and the given point is on the terminal side. Then find the values of the 6 trig functions.

a. $(-3, -4)$

b. $(7, 4)$

c. $(-\sqrt{2}, \sqrt{2})$

Examples: Finding the 6 Trig Functions Given an Equation of the Terminal Side

An equation of the terminal side of an angle θ (in standard position) is given with a restriction on x . Sketch the situation with least positive angle θ , and find the values of the 6 trig functions.

a. $-6x - y = 0, x \geq 0$.

b. $x + y = 0, x \geq 0$

c. $x = 0, y \geq 0$.

Lesson: Quadrantal Angles and Trig Functions

What were the quadrantal angles?

If one of the quadrantal angles is involved, how does this impact how we think about the trig functions? Draw a sketch to help illustrate this point. Explain how these are different from non-quadrantal angles, and how we choose points in these cases.

Example Find the indicated value of: $\sin 90^\circ$.

How should we think about something like $\sin^2 90^\circ$?

Example Use trigonometric function values of quadrantal angles to evaluate: $\tan 0^\circ - 6 \sin 90^\circ$.

Example Use trigonometric function values of quadrantal angles to evaluate: $\sin^2 180^\circ + \cos^2 180^\circ$.

Examples: Quadrantal Angles and Trig Functions

Find the function values:

a. $\cot 90^\circ$

b. $\sec 180^\circ$

c. $\sin (-270^\circ)$

d. $3 \sec 180^\circ - 2 \tan 360^\circ$

e. $\sin^2 90^\circ - 3 \cos^2 90^\circ$

Lesson: Reference Angles and Non-Acute Angles

Review: what is a quadrantal angle?

What is associated with every non-quadrantal angle?

Show how we can find reference angles by drawing the reference graphs in the video:

Example Find the reference angle for each angle.

a. 274°

b. 822°

Draw how to find the reference angle based on the quadrant (follow along with the video and draw the diagram on your own):

Now let's talk about using 30° , 45° , and 60° as reference angles.

Example Find the values of the 6 trig functions for 240° . Draw a reference!!! (This class is all about pictures!!)

Don't forget the helpful device: All Students Take Calculus 😊

Let's outline the steps for finding trig function values for any non-quadrantal angle:

1.

2.

3.

4.

How can you avoid making an error with finding a trig function value?

Examples Find the exact values:

a. $\sin(-150^\circ)$

b. $\cos(-510^\circ)$

Example Evaluate: $\sin^2 120^\circ + \cos^2 120^\circ$

Example Find all values of θ , if θ is in the interval $[0^\circ, 360^\circ)$ and has the given function value: $\cos \theta = \frac{\sqrt{3}}{2}$

Example Is the following true or false? $\cos 60^\circ = 2 \cos 30^\circ$

Examples: Reference Angles and Finding Trig Value Functions

Find the exact values of all 6 trig functions of each angle. Rationalize when applicable.

a. 315°

b. -855°

Evaluate

a. $\cot^2 120^\circ + \tan^2 60^\circ - \cos^2 330^\circ$

b. $\cot^2 135^\circ - \tan^4 45^\circ - \sin^2 150^\circ$

Examples: Finding the Value of Theta

Find all values of θ if θ is in the interval $[0^\circ, 360^\circ)$ and has the given value.

a. $\sec \theta = -\sqrt{2}$

b. $\cot \theta = -\frac{\sqrt{3}}{3}$

c. $\cos \theta = \frac{1}{2}$

Lesson: Sine and Cosine as Periodic Functions

What is a periodic function?

What are some examples?

What are 2 known periodic functions?


What can we say about the behavior of these functions as we move along the unit circle?

AS s CHANGES FROM	THE VALUE OF $\sin s$	THE VALUE OF $\cos s$
0 TO $\frac{\pi}{2}$		
$\frac{\pi}{2}$ TO π		
π TO $\frac{3\pi}{2}$		
$\frac{3\pi}{2}$ TO 2π		

But let's change s to x going forward so we don't get confused.

Draw the graph of sine, filling in the table and then using the graph paper.


x	$y = \sin x$
0	
$\frac{\pi}{4}$	
$\frac{\pi}{2}$	
π	
$\frac{3\pi}{2}$	
2π	



 Math-Aids.Com

Draw the graph of cosine, filling in the table and then using the graph paper.

x	$y = \sin x$
0	
$\frac{\pi}{4}$	
$\frac{\pi}{2}$	
π	
$\frac{3\pi}{2}$	
2π	



 Math-Aids.Com

What term do we use to describe the graph of sine?

How do we describe the graph of cosine?

What is the period of both the sine and cosine functions?

Where are the graphs continuous?

What is the domain and range of the graphs?

How can we describe the x-intercepts of sine?

How can we describe the x-intercepts of cosine?

What is the test for even or odd functions?

What does it mean to be an even function?

What does it mean to be an odd function?

How do you know that $y = \sin x$ is odd? How would you show it?

How do you know that $y = \cos x$ is even? How would you show it?

Lesson: Using Amplitude and Period to Graph Sine and Cosine

What is amplitude?

Using the equation $y = a \sin x$, when will this graph stretch or shrink? Draw 2 comparison sketches to help you remember (just a quick sketch so you remember how these visually compare)

Using the equation $y = a \cos x$, when will this graph stretch or shrink? Draw 2 comparison sketches to help you remember (just a quick sketch so you remember how these visually compare)

What happens when a is negative? Demonstrate this with some brief sketches so that you can remember.

What is period?

Using the equation $y = \sin bx$, watch the video and write an explanation with sketches to demonstrate period.

Using the equation $y = \cos bx$, watch the video and write an explanation with sketches to demonstrate period.

What formula can you use to determine the period of a function?

Examples Determine the periods of the following functions:

a. $y = \sin 3x$

b. $y = \cos \frac{1}{2}x$

c. $y = \sin 2\pi x$

Summarize amplitude and period when the function is of the form $y = a \sin bx$ and $y = a \cos bx$

Examples Graph the following over 2 period intervals, completing the table with strategic points to help you graph. Give the period and amplitude in each. You will have to sketch your own graph and create your own axes.

$$y = \cos \frac{1}{3}x$$

Period:


Amplitude:

x	$y = \cos \frac{1}{3}x$

$$y = 2 \sin x$$

Period:

Amplitude:



$$y = 3 \cos 2\pi x$$

Period:

Amplitude:

x	y

Examples: Using Amplitude and Period to Graph Sine and Cosine

Graph the following over two-period intervals.

a. $y = -\frac{1}{2}\sin x$

b. $y = 3 \cos 2x$


d. $y = -\sin \pi x$

Lesson: Determining the Equations of Sine and Cosine from Graphs


In each graph, determine the equation of the form $y = a \cos bx$ or $y = a \sin bx$, with $b > 0$, for each graph.

Desmos.com was used to create these graphs.


a.




b.



c.



d.



Lesson: Translations of the Sine and Cosine Graphs

How do you describe a horizontal translation? In your own words / explanation, summarize with sketches how the video show these translations work.

What is phase shift?

What is an argument?

How do you describe a vertical translation? In your own words / explanation, summarize with sketches how the video show these translations work.

What happens when a function is of the form: $f(x) = a\cos(bx + bd)$ with the phase shift?

Examples Identify the amplitude, period, phase shift, argument, and vertical translation.

a. $y = \sin\left(x + \frac{\pi}{4}\right)$

e. $y = 2 + \sin(2x - 2\pi)$

b. $y = 2 - \sin x$

f. $y = 1 - 2 \cos\left(3x + \frac{\pi}{2}\right)$

c. $y = 3 + \sin(x + 2)$

g. $y = 2 - 4 \cos(3x + 1)$

d. $y = -4 + \frac{1}{2} \cos x$

h. $y = 3\pi - 2 \sin(\pi x - 3)$

Examples Show how to find the start and end points of an interval for us to graph.

a. $y = \sin\left(x + \frac{\pi}{2}\right)$

b. $y = 1 - 2 \sin\left(3x - \frac{\pi}{2}\right)$

c. $y = 3\pi - 2 \sin(\pi x - 3)$

Example Using the graph $y = \sin(x + \frac{\pi}{4})$, and *clearly write out the steps* required to graph any sine or cosine graph. Identify the amplitude, period, phase shift, and vertical translation.

Example Graph $y = 1 + \cos x$. Identify the amplitude, period, phase shift, and vertical translation.

Example Graph $y = -1 + \frac{1}{2}(\cos 2x - \pi)$. Identify the amplitude, period, phase shift, and vertical translation.

Examples: Translations with Sine and Cosine


Graph the following: $y = \frac{5}{2} \sin\left(4\left(x + \frac{\pi}{4}\right)\right)$

Lesson: The Graphs of Tangent and Cotangent

What is an asymptote? Draw a sketch to help explain.


Draw the graph of tangent, filling in the table and then using the graph paper.

x	$y = \tan x$
$-\frac{\pi}{2}$	
$-\frac{\pi}{4}$	
0	
$\frac{\pi}{4}$	
$\frac{\pi}{2}$	
$\frac{3\pi}{2}$	



Draw the graph of cotangent, filling in the table and then using the graph paper.

x	$y = \cot x$
0	
$\frac{\pi}{4}$	
$\frac{\pi}{2}$	
$\frac{3\pi}{2}$	
π	
$\frac{5\pi}{4}$	



 Math-Aids.Com

How would we describe the period for these functions?

Where are the graphs continuous (and discontinuous)?

What is the domain and range of the graphs?

How can we describe the x-intercepts of tangent?

How can we describe the x-intercepts of cotangent?

Are either function even or odd?

How do you interpret amplitude with these functions and why?

What is the key for graphing? (Be specific with the equations you need to leverage)

Example Graph over a one-period interval: $y = \tan 2x$

Example Graph over a two-period interval: $y = -\frac{1}{2} \cot 4x$

Lesson: Translations with The Graphs of Tangent and Cotangent

Tangent and cotangent can experience the same types of translations that sine and cosine did (which we discussed in a previous lesson).

Examples Describe the period, phase shift, and vertical translations for each function.

a. $y = 3 \tan 4x$

d. $y = 3 + \cot(x - 4)$

b. $y = -\cot(x + \pi)$

e. $y = -2 + \tan\left(2\pi x - \frac{\pi}{4}\right)$

c. $y = 2 \tan\left(3x - \frac{\pi}{2}\right)$

f. $y = \frac{1}{2} - \cot(5x + 2)$

Examples Show how to find 2 adjacent asymptotes for graphing each of the following:

a. $y = 3 \tan 4x$

e. $y = -2 + \tan\left(2\pi x - \frac{\pi}{4}\right)$

b. $y = 2 \tan\left(3x - \frac{\pi}{2}\right)$

f. $y = \frac{1}{2} - \cot(5x + 2)$

Example Graph $y = 3 + \cot\left(x + \frac{\pi}{4}\right)$

Example Graph $y = -1 + 2\tan(4x + \pi)$

Examples: Translations with The Graphs of Tangent and Cotangent

Graph over a two-period interval.

a. $y = \tan\left(3x + \frac{\pi}{4}\right)$


b. $y = -2 + 3 \cot(4x - \pi)$

c. $y = 1 - \tan(2\pi x - \pi)$


Lesson: Determining the Equations of Tangent and Cotangent from Graphs

Graphs from desmos.com of course!


a.



b.




C.



Lesson: Translations of The Graphs of Secant and Cosecant

Draw the graph of secant, filling in the table and then using the graph paper.


x	$y = \sec x$
$-\frac{\pi}{2}$	
$-\frac{\pi}{4}$	
0	
$\frac{\pi}{4}$	
$\frac{\pi}{2}$	
$\frac{3\pi}{4}$	
π	
$\frac{3\pi}{2}$	



 Math-Aids.Com

Draw the graph of cosecant, filling in the table and then using the graph paper.

x	$y = \sec x$
0	
$\frac{\pi}{6}$	
$\frac{\pi}{3}$	
$\frac{\pi}{2}$	
$\frac{2\pi}{3}$	
π	
$\frac{3\pi}{2}$	
2π	



 Math-Aids.Com

How would we describe the period for these functions?

Where are the graphs continuous (and discontinuous)?

What is the domain and range of the graphs?

How can we describe the x-intercepts of both graphs?

Are either function even or odd?

How do you interpret amplitude with these functions?

What is the key for graphing? (Be specific with the equations you need to leverage)

Examples Graph $y = 2 \csc \frac{1}{2}x$

Examples Graph $y = 1 + \frac{1}{2}\sec\left(x - \frac{\pi}{2}\right)$

Examples: Translations of The Graphs of Secant and Cosecant

Graph the following over a one-period interval.

a. $y = -2 \sec\left(x + \frac{3\pi}{4}\right)$

b. $y = \frac{1}{2} \csc(2x - \pi)$


c. $y = 2 + \csc\left(\frac{1}{2}x - \pi\right)$

d. $y = 2 - 3 \sec\left(2\pi x - \frac{\pi}{2}\right)$


Lesson: Equations of Secant and Cosecant from Graphs

Determine the equation of secant or cosecant from the graphs.


a.



b.



C.



Lesson: Inverse Functions (a review)

What does it mean to be an “inverse function” ?

Example If a function includes the points $\{(2,3), (5,4), (-7,2), (-1,5)\}$, then what points will the inverse have?

What is the horizontal line test?

Example Draw the graphs from the video, then determine if they have inverses:

How do you find the inverse of a function $f(x)$?


Example Find the inverse of the following:

a. $f(x) = 2x + 1$


b. $f(x) = \sqrt{x}$

Examples Graph the following functions with their inverses:

a. $f(x) = 2x + 1$



b. $f(x) = \sqrt{x}$



From these graphs, we can see that inverses of functions are really just what?

What can we say about the domain and range inverses of functions?

Lesson: Inverse Trig Functions

Does the sine function have an inverse? Why or why not?

How can we manipulate the graph of the sine function to determine an inverse?

What are two ways that we denote the inverse sine function?

How do we “formally” define the inverse sine function?

Show how to graph the inverse sine function. Draw both the table and graph discussed from the video:

Example Find the value of each of the following:

a. $y = \arcsin \frac{\sqrt{3}}{2}$

b. $f(x) = \sin^{-1} \left(-\frac{\pi}{4} \right)$

c. $y = \sin^{-1} -\sqrt{2}$

Does the cosine function have an inverse? How should we think about this?

How do we denote the inverse of cosine?

How do we define the inverse cosine function?

Show how to graph the inverse cosine function. Draw both the table and graph discussed from the video:

Examples Find the value of each of the following:

a. $y = \arccos 1$

b. $f(x) = \cos^{-1}\left(\frac{1}{2}\right)$

How do we denote the inverse of tangent?

How do we define the inverse tangent function?

Show how to graph the inverse tangent function. Draw both the table and graph discussed from the video:

Define the other inverse circular functions:

Summarize the domains, ranges, and quadrants of the inverse trig functions below (this is important, you'll need to study this for the next exam):

Function	Domain	Interval	Quadrants

Examples Find the degree measure of θ :

a. $\theta = \arctan \sqrt{3}$

b. $\theta = \csc^{-1}(-\sqrt{2})$

Lesson: Finding Values of Inverse Trig Functions

Here we will just examine a wealth of examples of how to find different values of inverse trig functions.

Examples Use a calculator to determine the appropriate values.

a. Find y in radians if $y = \sec^{-1}(-4)$

b. Find θ in degrees if $\theta = \operatorname{arccot}(-0.25431)$

Before you evaluate those, you will have to watch some explanation of how / why we evaluate these the way that we do in the calculator. Write out the explanation of how/why we evaluate $\csc^{-1}(2)$ in a calculator below:

Cheat Sheet – How to Evaluate Inverse Trig Functions That Don't Have a Button on a Calculator

Cheat Sheet (for Comparison) – How to Evaluate Regular Trig Functions That Don't Have a Button on a Calculator

This might get confusing so maybe rip this sheet out and put it someplace where you can easily reference it??

Examples Use a calculator to find each value.

a. $\sin(\cos^{-1} 0.25)$

b. $\cot(\cos^{-1}(0.56331233))$

Examples Evaluate each expression without using a calculator:

a. $\cos\left(\sin^{-1}\left(\frac{2}{3}\right)\right)$

b. $\sec(\sec^{-1} 2)$

c. $\sin\left(\tan^{-1}\frac{4}{3} - \cos^{-1}\frac{12}{13}\right)$

Examples Write each expression as an algebraic expression in terms of u .

a. $\cot(\sec^{-1} u)$, $u > 0$

b. $\sec\left(\operatorname{arccot}\left(\frac{\sqrt{4-u^2}}{u}\right)\right)$

Lesson: Reciprocal Identities of Trig Functions

First, what is a reciprocal? Give an example.

What are the 6 reciprocal identities?

1.

2.

3.

4.

5.

6.

You should NOT write out something like _____ or _____ as a shortcut for a reciprocal identities. Why not?

How can you shortcut the reciprocal identity then with notation?

What are some other ways we can relate the reciprocal identities to one another?

Examples Find the function values.

a. $\tan \theta$ given that $\cot \theta = 3$

b. $\sec \theta$ given that $\cos \theta = -\frac{5}{\sqrt{50}}$

c. $\sin \theta$ given that $\csc \theta = 1.25$

Lesson: Signs and Ranges of Trig Functions

Review: what does r represent when we are talking about trig functions?

Since r is a distance, we know _____.

What determines if a trig function has a positive or negative value?

Draw a picture to help breakdown and analyze the signs of theta:

Now fill in this table to help you remember the sign of each function:

θ IN QUADRANT	$\sin \theta$	$\cos \theta$	$\tan \theta$	$\cot \theta$	$\sec \theta$	$\csc \theta$
I						
II						
III						
IV						

One way to remember the signs is with the phrase All Students Take Calculus. Break down this this is helpful.

Examples Determine the signs of the trig functions of an angle in standard position with the given measure.

a. 23°

b. 255°

c. -30°

Examples Determine which quadrant of an angle θ that satisfies the given conditions.

a. $\sin \theta > 0, \csc \theta > 0$

b. $\sec \theta > 0, \csc \theta > 0$

Now we will show how to get ranges of the six trig functions.

Draw a picture to demonstrate how angle θ increase in measure from $0^\circ - 90^\circ$.

As the measure of an angle _____, _____ but never exceeds _____. Therefore:

Let's repeat this idea but in quadrant IV. Follow along with the video below and write down the observations again.

What can we conclude from these sketches and observations?

What can we conclude about the values of $\tan \theta$ and $\cot \theta$? And why?

What can we conclude about the values of $\sec \theta$ and $\csc \theta$? Why?

We have now discovered the ranges of all the trig functions. Let's summarize this in the table.

TRIG FUNCTION	RANGE (WRITE IN BOTH SET-BUILDER NOTATION AND INTERVAL NOTATION)
$\sin \theta, \cos \theta$	
$\tan \theta, \cot \theta$	
$\sec \theta, \csc \theta$	

Example Determine if each statement is possible or impossible.

a. $\cot \theta = -0.98$

b. $\cos \theta = -1.25$

c. $\sec \theta = 0.72$

Lesson: Pythagorean and Quotient Identities

Show how we can build the first Pythagorean identity. These are actually pretty darn important, so you really do need to understand this argument!!

Now let's build another one:

And now let's build the last one:

Summarize the Pythagorean identities:

What is only key about these identities?

Now let's show how to build the quotient identities:

By the way, similar reasoning holds for $\cot \theta$.

Summarize the Quotient Identities below:

Example Find the value of $\cos \theta$ and $\tan \theta$, given that $\sin \theta = -\frac{\sqrt{2}}{2}$ and $\cos \theta > 0$.

Example Find $\sin \theta$, given that $\cos \theta = \frac{4}{5}$ and θ is in Quadrant IV.

Example Give all 6 trig function values given that $\cot \theta = \frac{\sqrt{3}}{8}$ and θ is in quadrant I.

Examples: Using Pythagorean and Quotient Identities to Find Trig Function Values

Find all 6 trig function values given the information.

a. $\cos \theta = -\frac{3}{5}$, θ is in quadrant III.

b. $\csc \theta = 2$, θ is in quadrant II.

c. $\cos \theta = 1$

Lesson: Fundamental Identities of Trig

Review the test for symmetry (and the results):

We know that the sine function is odd, show what this means:

We know that the cosine function is even, show what this means:

These are called **identities**.

Using these identities, what other identity can we determine about the tangent function:

These are called the **even-odd identities**.

We have already seen a handful of identities, and we call these the fundamental identities. Summarize them below.

Reciprocal Identities

Quotient Identities

Pythagorean Identities

Even-Odd Identities

Example If $\cos \theta = \frac{3}{4}$ and θ is in quadrant I, find $\sin \theta$.

Example If $\tan \theta = -\frac{1}{4}$ and θ is in quadrant IV, find the values of all the other 5 trig functions.

We use the identities to work on problems called transformations.

Example Write $\tan x$ in terms of $\sec x$.

Example Write the following in terms of sine and cosine, and then simplify until no quotients appear and all functions are of θ only.

a. $\csc \theta \cos \theta \tan \theta$

b. $\cos \theta (\cos \theta - \sec \theta)$

Examples: Using Fundamental Trig Identities to Find Values of Other Trig Functions

Find the five remaining trig functions given the following information.

a. $\csc \theta = -\frac{5}{2}$, θ in quadrant III.

b. $\sec \theta = \frac{7}{2}$, $\tan \theta < 0$

Examples: Writing Trig Functions in Terms of Other Trig Functions (Fundamental Identities)

a. Write $\cot x$ in terms of $\sin x$

b. Write $\csc x$ in terms of $\cos x$

Examples: Using Fundamental Identities to Write Expressions in Terms of Sine and Cosine

Write each expression in terms of sine and cosine, then simplify so that no quotients appear and all functions are in θ only.

a. $\cos \theta \csc \theta$

b. $\frac{(1+\cot \theta)}{\cot \theta}$

c. $\frac{\sec^2 \theta - 1}{\csc^2 \theta - 1}$

d. $\frac{\csc \theta}{\cot(-\theta)}$

Lesson: Verifying Trigonometric Identities

As you progress through higher mathematics, sometimes you will need to manipulate equations in very creative ways. One way to learn this level of creativity is by verifying trigonometric identities. We will go through a few techniques of how to do this.

Strategy 1: Rewrite the More Complicated Side / Work from One Side

a. Verify that $\frac{1-\sin^2 x}{\cos x} = \cos x$. Show *why*.

b. Verify that $\sin^2 x + \tan^2 x + \cos^2 x = \sec^2 x$

Strategy 2: Express the Functions in Terms of All Sine and Cosine

a. $\frac{\tan x}{\sec x} = \sin x$

b. $\sec x (\sin x + \cos x) = 1 + \tan x$

Strategy 3: Factor or Complete Indicated Algebraic Operations

Examples Factor.

a. $x^2 - 4$

b. $\sin^2 x - 4$

c. $(\tan x + \cot x)^2 - (\tan x - \cot x)^2$

d. $2 \sin^2 x + 3 \sin x + 1$

Examples Verify the following identities.

a. $\sin^4 x - \cos^4 x = 2 \sin^2 x - 1$

b. $\frac{1-\cos x}{1+\cos x} = (\cot x - \csc x)^2$

Strategy 4: Manipulate Both Sides

a. $\frac{\cos x}{\sec x} + \frac{\sin x}{\csc x} = \sec^2 x - \tan^2 x$

b. $\tan^2 x \sin^2 x = \tan^2 x + \cos^2 x - 1$

Strategy 5: Use the Conjugate

What is the conjugate of the following:

a. $x - 4$

b. $\sqrt{2} - \sqrt{3}$

c. $\sin x + 1$

Verify the following identities: $\frac{1-\cos x}{1+\cos x} = \csc^2 x - 2 \csc x \cot x + \cot^2 x$

Examples: Verifying Trigonometric Identities

a. $\frac{\cos x}{\cot x \sin x} = 1$

b. $\frac{1}{\sec x - \tan x} = \sec x + \tan x$

c. $\frac{1+\sin t}{1-\sin t} - \frac{1-\sin t}{1+\sin t} = 4 \tan t \sec t$

d. $(\cos \theta - \sin \theta)(\sec \theta + \csc \theta) = \cot \theta - \tan \theta$

Lesson: Proofs of the Sum and Difference Identities for Cosine

Write out the proof for $\cos(A - B) = \cos A \cos B + \sin A \sin B$. Do your best to understand the proof and then explain it in a way that you will understand.

Write out the identity for $\cos(A - B)$ below (yes I know I stated it above, but YOU should write it so you have a better chance of remembering it):

Write out the proof for $\cos(A + B) = \cos A \cos B - \sin A \sin B$

Write out the identity below (yes, write it out so you don't forget!):

Lesson: Using Sum and Difference Identities for Cosine

Examples Show how to use the cosine of a sum or difference to find the exact value of the expressions:

a. $\cos 105^\circ$

b. $\cos \frac{\pi}{12}$

c. $\cos 40^\circ \cos 50^\circ - \sin 40^\circ \sin 50^\circ$

Show how the cofunction identities are an extension of the cosine of a difference:

Write out all the **cofunction identities**

Examples Find one value of θ that satisfies the following:

a. $\sec \theta = \csc 72^\circ$

b. $\tan \theta = \cot(-44^\circ)$

c. $\cos \theta = \sin \frac{5\pi}{6}$

Example Write $\cos(180^\circ - \theta)$ as a trigonometric function of θ alone.

Example Suppose $\sin s = \frac{3}{5}$ and $\sin t = -\frac{12}{13}$, where s is in quadrant I and t is in quadrant III. Find $\cos(s + t)$ and $\cos(s - t)$.

Example Verify that $\sec(\pi + x) = -\sec x$

Lesson: Sum and Difference Identities for Cosine

Write out the proof for $\sin(A + B)$

Write out the proof for $\sin(A - B)$

Write out the identities for the sum and difference of sine below:

Write out the proof for $\tan(A + B)$

Write all identities for the sum and difference of tangent below:

Examples Find the exact value of each expression.

a. $\sin(-15^\circ)$

b. $\tan\left(-\frac{5\pi}{12}\right)$

c. $\sin\left(\frac{5\pi}{9}\right)\cos\left(\frac{\pi}{18}\right) + \cos\left(\frac{5\pi}{9}\right)\sin\left(\frac{\pi}{18}\right)$

Examples Write each function as an expression involving functions of θ alone.

a. $\cos(30^\circ + \theta)$

b. $\sin(45^\circ + \theta)$

c. $\tan\left(\frac{\pi}{4} + x\right)$

Examples Suppose A and B are angles in standard position such that $\sin A = -\frac{3}{5}$, $\frac{3\pi}{2} < A < 2\pi$ and

$\cos B = -\frac{7}{25}$, $\pi < B < \frac{3\pi}{2}$. Find the following.

a. $\sin(A + B)$

b. $\tan(A - B)$

c. the quadrant of A-B

Examples Verify the identities.

a. $\sin(x + y) + \sin(x - y) = 2 \sin x \cos y$

b. $\frac{\sin(x-y)}{\sin(x+y)} = \frac{\tan x - \tan y}{\tan x + \tan y}$

Examples: Sine of a Sum and Tangent of a Sum

Let $\cos s = \frac{5}{12}$, $\sin t = \frac{4}{5}$ where s and t are in quadrant I. Find:

$$\sin(s + t)$$

$$\tan(s + t)$$

The quadrant of $(s + t)$

Lesson: Double-Angle Identities

Show how we derive the first identity for $\cos 2A$:

Show how we derive the second identity for $\cos 2A$:

Show how we derive the third identity for $\cos 2A$:

Show how we derive the identity for $\sin 2A$:

Show how we derive the identity for $\tan 2A$:

Summarize the Double-Angle Identities Below:

Example Give $\sin \theta = \frac{5}{8}$ and $\cos \theta < 0$, find $\sin 2\theta$, $\cos 2\theta$, and $\tan 2\theta$.

Example Find the values of the six trig functions give $\cos 2\theta = -\frac{11}{13}$ and $180^\circ < \theta < 270^\circ$.

Example Verify the following identities:

a. $(\sin x + \cos x)^2 = \sin 2x + 1$

b. $\frac{\cot x - \tan x}{\cot x + \tan x} = \cos 2x$

Examples Simplify the expressions.

a. $\sin^2 15^\circ - \cos^2 15^\circ$

b. $\sin^2 \frac{3\pi}{5} - \cos^2 \frac{3\pi}{5}$

Example Write $\cos 3x$ in terms of $\cos x$

Lesson: Product-to-Sum Identities

Show what happens when we add the identities $\cos(A + B)$ to $\cos(A - B)$

Now show what happens when we subtract them:

Summarize the Product-to-Sum and Sum-to-Product Identities

Examples Write the following as the sum or difference of 2 trig functions.

a. $2 \sin 48^\circ \cos 104^\circ$

b. $8 \sin 9x \sin 5x$

Examples Write the expression as a product of trigonometric functions.

a. $\cos 6x + \cos 9x$

b. $\sin 106^\circ - \sin 96^\circ$

Lesson: Half-Angle Identities

Show how we derive the identity for $\sin \frac{A}{2}$:

Show how we derive the identity for $\cos \frac{A}{2}$:

Show the first way we can derive an identity for $\tan \frac{A}{2}$:

Show the second way we can derive an identity for $\tan \frac{A}{2}$:

Show the third way we can derive an identity for $\tan \frac{A}{2}$:

Summarize the half-angle identities below:

Example Show how to use the half-angle identity to evaluate: $\sin 12.5^\circ$

Example Find the exact value of $\cos 165^\circ$

Example Find the exact value of $\tan 75^\circ$ using the identity $\tan A = \frac{\sin A}{1 + \cos A}$

Example Given $\cos s = \frac{2}{7}$ with $0 < s < \frac{\pi}{2}$, find $\sin \frac{s}{2}$, $\cos \frac{s}{2}$, and $\tan \frac{s}{2}$

Example Simplify each expression.

a. $\sqrt{\frac{1 - \cos 40^\circ}{2}}$

b. $\frac{1 - \cos 64.72^\circ}{\sin(64.72^\circ)}$

c. $\pm \sqrt{\frac{1 + \cos 10x}{2}}$

Example Verify the identities.

a. $\cot^2 \frac{x}{2} = \frac{(1+\cos x)^2}{\sin^2 x}$

b. $\cos x = \frac{1-\tan^2 \frac{x}{2}}{1+\tan^2 \frac{x}{2}}$

Lesson: Summarizing All of the Identities

Using all the trig identities that we've discussed, verify the following:

a. $\frac{\sin x}{1+\cos x} = \frac{1-\cos x}{\sin x}$

b. $1 - \tan^2 \frac{x}{2} = \frac{2 \cos x}{1+\cos x}$

c. $\tan\left(\frac{a}{2} + \frac{\pi}{4}\right) = \sec a + \tan a$

d. $\sin 7x + \sin 5x + \sin 3x + \sin x = 4 \sin 4x \cos 2x \cos x$

$$\text{e. } \tan 4x = \frac{2 \tan 2x}{2 - \sec^2 2x}$$

Lesson: Solving Trig Equations, part 1 – linear methods

Recall how linear methods work first by solving: $2x + 4 = 8$

Now compare this approach with the equation: $3 \tan \theta - \sqrt{3} = 0$, over the interval $[0^\circ, 360^\circ)$

How does this answer change if we want to solve for *all* solutions?

Example Solve: $2 \sin x + 3 = 4$ over $[0, 2\pi)$

Lesson: Solving Trig Equations, part 2 – quadratic methods

Using quadratic methods to solve trig equations relies on using the zero-factor property. Show how the zero-factor property works by solving $(x + 5)(x - 3) = 0$.

Compare this approach by solving $\cos \theta \sin \theta = 0 = -\cos \theta$, over $[0^\circ, 360^\circ)$.

Example Solve $(\csc x + 2)(\csc x - \sqrt{2}) = 0$ for all solutions.

Recall how to use quadratic methods by solving $x^2 + 5x + 6 = 0$.

Compare this approach by solving $\cos^2 x + 2 \cos x + 1 = 0$ over $[0^\circ, 360^\circ)$.

Example Solve $2 \cos^2 x + \cos x - 1 = 0$ for all possible solutions.

Example Solve $2 \cos^2 x - \sqrt{3} \cos x = 0$ over $[0, 2\pi)$

Example Solve $\tan x \csc x - \sqrt{3} \csc x = 0$ for all solutions.

Example Solve $\csc^2 x - 2 \cot x = 0$ over $[0^\circ, 360^\circ)$

Example Solve $\tan x - \sqrt{3} = \sqrt{3} \cot x - 1$ for all possible solutions.

Lesson: Solving Trig Equations, part 3 – equations with half-angles

Example Solve $2 \cos \frac{x}{2} - \sqrt{2} = 0$ over the interval $[0, 2\pi)$

How would the answers change if you needed to find all solutions?

Example Solve $2\sqrt{3}\sin\frac{x}{2} = 3$ for all possible solutions.

Example Solve $\sin^2\frac{x}{2} - 2 = 0$ for all possible solutions.

Lesson: Solving Trig Equations, part 4 – equations with multiple angles

Example Solve $\cos 2x = \sin x$ over $[0, 2\pi)$

Example Solve $2 \cos^2 x - 2 \sin^2 x + 1 = 0$ over $[0^\circ, 360^\circ)$

How would this change for all solutions?

Example Solve $\cot 2x - \csc 2x = 1$ over $[0, 2\pi)$

Example Solve $\tan 2x - \sec 2x = 2$ over $[0, 2\pi)$

Example Solve $\cos x - 1 = \cos 2x$ for all solutions.

Example Solve $\sin 2x = 2 \cos^2 x$ for all solutions.

Lesson: Equations Involving Inverse Trig Functions

Example Solve $y = 4 \tan 3x$, where x is restricted to $\left(-\frac{\pi}{6}, \frac{\pi}{6}\right)$.

Example Solve $y = \tan (2x - 1)$ for x in $\left(\frac{1}{2} - \frac{\pi}{4}, \frac{1}{2} + \frac{\pi}{4}\right)$

Example Solve $3 \tan^{-1} x = \pi$

Example Solve $6 \arccos\left(x - \frac{\pi}{3}\right) = \pi$

Example Solve $\cot^{-1} x = \tan^{-1} \frac{4}{3}$

Example Solve $\arccos x + 2 \arcsin \frac{\sqrt{3}}{2} = \frac{\pi}{3}$

Example Solve $\sin^{-1} x + \tan^{-1} x = 0$

Lesson: Oblique Triangles and the Law of Sines

First, we should review some facts about congruent triangles. When are 2 triangles congruent?

Congruence Axioms

Side-Angle-Side (SAS)

Angle-Side-Angle (ASA)

Side-Side-Side (SSS)

When does Side-Angle-Angle (SAA) hold?

When a triangle is not a right triangle, what is it called?

When can a triangle be solved?

What data is required for solving oblique triangles?

Case 1:

Case 2:

Case 3:

Case 4:

Is there an Angle-Angle-Angle congruency?

When Case 1 or Case 2 is involved, what is required?

When Case 3 or Case 4 is involved, what is required?

This Law of Cosines will be discussed in another video.

Show how we derive the Law of Sines:

State the Law of Sines:

What is an alternative form of the law of sines?

Example Solve Triangle ABC if $A = 28.5^\circ$, $C = 102.3^\circ$, $c = 25.1$ in.

Recall the formula for the area of a triangle:

Using SAS, how else can you find the area of a triangle?

Example Reference the sketch in the video. Find the area of the triangle.

Example Find the area of a triangle DEF if $D = 58^\circ 10'$, $e = 31.2in$, $F = 73^\circ 30'$.

Lesson: Applications of the Law of Sines

Example An engineer wishes to measure the distance across a river. Reference the sketch in the video.

If $C = 117.4^\circ$, $A = 30.0^\circ$, $b = 75.8$ ft

Review the two methods of bearing:

Example The bearing of a lighthouse from a ship was found to be $N 52^\circ W$. The ship sailed 12.3 miles south, and the new bearing was $N 23^\circ W$. Find the distance, to the nearest mile, between the ship and the lighthouse at each location.

Example A balloonist is directly above a straight road, 2 miles long that joins the 2 villages. The closer town to him is an angle of depression of 45° , and the farther town is at angle of depression of 41° . How high above the ground is the balloon?

Lesson: The Ambiguous Case of the Law of Sines

Describe the ambiguous case of the law of sines as describe in the video. There's going to be quite a bit of information presented – watch the entire explanation of why we need the law of sines and do your best to describe the various cases to sum up. Summarize the explanation so that it makes sense to YOU.

What were some tips for applying the law of sines?

Example Solve triangle ABC if $a = 16.2\text{ cm}$, $c = 14.1\text{ cm}$, and $C = 72^\circ 30'$.

Example Solve triangle DEF if $D = 62^\circ 8'$, $a = 34.5in$, $b = 38.1in$.

Explain the number of triangles satisfying the ambiguous case (SSA):

Examples Outline the steps to solve these triangles. You don't have to solve completely, but do enough work to determine how many possible triangles there are. Include sketches.

a. $B = 80^\circ, C = 35^\circ, b = 27$

b. $A = 60^\circ, B = 70^\circ, c = 5$

Examples Outline the steps to solve these triangles. You don't have to solve completely, but do enough work to determine how many possible triangles there are. Include sketches.

c. $a = 15, b = 24, B = 33^\circ$

d. $A = 25^\circ, b = 8, a = 11$

Examples Outline the steps to solve these triangles. You don't have to solve completely, but do enough work to determine how many possible triangles there are. Include sketches.

e. $A = 61.4^\circ, b = 35.5, b = 39.1$

f. $a = 17.1, c = 13.1, C = 76^\circ$

Example Without using the law of sines, explain why no triangle ABC exists satisfying $B = 95^\circ, b = 40\text{cm}, c = 46\text{cm}$.

Lesson: The Law of Cosines

What is the property of triangle side length restriction?

Show how to derive the law of cosines:

State the Law of Cosines:

Where is one place where we've seen this?

Example Two boats leave a harbor at the same time, traveling on courses that make an angle of $81^{\circ}30'$ between them. When the slower boat has traveled 65.2 miles, the faster boat has traveled 84.3 miles. At that time, what is the distance between the boats?

Example Solve triangle ABC if $B = 72.4^\circ$, $a = 31.4ft$, $c = 44.3ft$.

Example Solve triangle DEF if $d = 24.2in$, $b = 42.6in$, $c = 59.1in$.

Example The simplest type of roof truss is a triangle. Refer to the sketch in the video. Find angle C to the nearest degree for the truss shown.

Explain the 4 Cases of Solving Oblique Triangles

Example: The Law of Cosines with SSS Triangles

Find the angles of the triangles given the following sides:

a. $a = 3ft, b = 5ft, c = 6ft$

b. $a = 324m, b = 421m, c = 298m$

Lesson: Heron's Formula for the Area of a Triangle

Explain how to use Heron's formula (as explained in the video):

Example Find the area of triangle ABC where $a = 12m$, $b = 16m$, $c = 25m$

Example Find the area of the Bermuda Triangle if the sides of the triangle have approximate lengths 800mi, 875mi, and 1250mi.


Lesson: Polar Equations and Graphs

There are other coordinate systems besides the rectangular coordinate system.


We will discuss the polar coordinate system!

What is a pole? What is the polar axis? Draw a sketch.

Here is what the polar coordinate system looks like:




It can also be in degrees:



How do we convert between rectangular and polar coordinates?


Examples Show how to plot each point, and then determine the rectangular coordinates of each point.

a. $P(3, 45^\circ)$




Examples Show how to plot each point, and then determine the rectangular coordinates of each point.

b. $P\left(-4, \frac{\pi}{3}\right)$



Examples Show how to plot each point, and then determine the rectangular coordinates of each point.

c. $R\left(3, -\frac{\pi}{2}\right)$



While a given point in the plane can only have 1 pair of rectangular coordinates, how many pairs of polar coordinates can it have?


Examples Determine the following:

a. Three other pairs of coordinates for the point $P(3, -110^\circ)$.


b. Two pairs of polar coordinates for the point with rectangular coordinates $\left(-\frac{\sqrt{3}}{2}, -\frac{1}{2}\right)$

What makes an equation a “polar equation” ?


Example Convert the equation to a polar equation, and then graph it: $y = x - 1$



Example Convert the equation to a polar equation, and then graph it: $x^2 + y^2 = 25$



Example Graph $r = 1 - \sin \theta$



Lesson: Trigonometric (Polar) Form of Complex Numbers

Complex numbers cannot be ordered like real numbers can. What is one way that we can organize complex numbers?

Show how we can graph complex numbers:

Complex numbers correspond to vectors. For a complex number $x + yi$, show how this corresponds to a vector (sketch), then label the sketch.

Examples Add $3 + 2i$ and $2 - 5i$, then graph both numbers and their resultant.

Define the relationships between x, y, r, θ below (with a sketch)

What is the trigonometric form (or polar form) of a complex number?

How do we refer to the number r ?

How do we refer to θ ?

Example Write $3(\cos 300^\circ + i \sin 300^\circ)$ in rectangular form.

How do you convert from rectangular form to trigonometric form?

Examples Write in trigonometric form.

a. $-\sqrt{2} + i$

b. -12

Examples Write the complex number in its alternative form.

a. $8(\cos 200^\circ + i \sin 200^\circ)$

b. $-8 + 5i$

Lesson: The Product and Quotient Theorems

First, let's review how to multiply $(2 + i\sqrt{3})(-2\sqrt{3} + 2i)$

Let's rewrite both of those complex numbers in their trigonometric form:

Now multiply their trigonometric forms together:

What do you notice about the result?

This problem demonstrates the product theorem.

State the Product Theorem:

Example Find the product of $3(\cos 45^\circ + i \sin 45^\circ)$ and $5(\cos 120^\circ + i \sin 120^\circ)$. Then write the answer in rectangular form.

(PART A) Show how to find the quotient of $(2\sqrt{6} - 2i\sqrt{2})$ and $(\sqrt{2} - i\sqrt{6})$

Now rewrite these complex numbers in their trigonometric form.

How can the “answer” to PART A be rewritten in trigonometric form?

What do you notice about this result?

This demonstrates the quotient theorem.

State the Quotient Theorem.

Example Find the quotient $\frac{35 \operatorname{cis} 45^\circ}{7 \operatorname{cis} (-90^\circ)}$. Write the result in rectangular form.

Example Use a calculator to find the following. Put your answer in rectangular form.

a. $(7.2 \operatorname{cis} 38.5^\circ)(3.5 \operatorname{cis} 49.8^\circ)$

b. $\frac{15.42 \left(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6}\right)}{5.21 \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4}\right)}$

Lesson: De Moivre's Theorem and Powers and Roots of Complex Numbers

Find the product: $[r (\cos \theta + i \sin \theta)]^2$

This is one example of De Moivre's theorem.

State De Moivre's Theorem:

Example Find $(\sqrt{3} + i\sqrt{5})^8$ and write the answer in rectangular form.

We can extend this theorem to find all n th roots of complex numbers.

Explain the concept of finding the n th root of a complex number.

State the n th Root Theorem

Example Find the three cube roots of -8

Example Find all complex number solutions to $x^4 + 1 = 0$.

Lesson: Vectors

What is a scalar?

What are vector quantities?

What is a vector?

Draw some vectors showing the differences with magnitude and directions.

Explain some of the notation we use regarding these concepts:

When are two vectors equal?

Is the sum of two vectors still a vector? (circle)

Yes

No

How do you find the sum of two vectors?

Vector addition is commutative. What does this mean?

For every vector, there is an opposite vector. Explain what this means:

The sum of $\mathbf{v}$ and $-\mathbf{v}$ results in what?

What happens when you multiply a vector and a scalar?

Explain the geometric properties of parallelograms:

Examples For each pair of vectors and the angle θ between them, sketch the result.

a. $|\mathbf{u}| = 12, |\mathbf{v}| = 22, \theta = 25^\circ$

b. $|\mathbf{u}| = 20, |\mathbf{v}| = 40, \theta = 30^\circ$

Examples Use the parallelogram in the video to find the magnitude of the resultant force for the 2 forces shown in the figure in the video.

Lesson: Vectors and Applications of Magnitude

Example A boat is being towed by two other boats, with the first boat having a force of 840lbs and the second have 925lb. The angle between these forces is 26° . Find the direction and magnitude of the equilibrant.

Example Two forces of 150 lbs and 275 lbs act at a point. The resultant force between them is 350 lbs. Find the angle between the forces.

Lesson: Vectors and Applications of Navigation

Example A ship leaves port on a bearing of 27° and travels 62.4 miles. The ship then turns due east and travels 85.2 miles. How far is the ship from the port? What is its bearing from port?

Lesson: Vectors and Applications of Incline

Example Find the force required to keep 2700-lb car parked on a hill that makes a 13° angle with the horizontal.

Example A force of 18.00 lb is required to hold a 74.0 lb crate on a ramp. What angle does the ramp make with the horizontal?

Lesson: Vectors (Algebraic Interpretation)

Vectors are defined algebraically in the rectangular coordinate system with what notation?

Draw how we interpret this vector below.

In a previous video, we discussed how to define vectors geometrically. How is this definition different from our algebraic interpretation?

Show how to define a direction angle.

How do we calculate the magnitude of a vector?

How do we calculate the direction angle of the same vector $\mathbf{u} = \langle a, b \rangle$?

Example Calculate the magnitude and direction for $\mathbf{u} = \langle 5, 2 \rangle$.

Draw how we can view the horizontal and vertical components of vectors.

Example A vector has magnitude 36.0 and direction angle 35° . Find the horizontal and vertical components.

Example Copy the sketch in the video. Write the vectors in the form $\langle a, b \rangle$.

What are the vector operations?

Example Let $\mathbf{u} = \langle -3, 4 \rangle$, $\mathbf{v} = \langle 5, 7 \rangle$. Find and illustrate each of the following.

a. $\mathbf{u} + \mathbf{v}$

b. $-3\mathbf{v} + 2\mathbf{u}$

What is the i, j form of vectors?

Example Write $\langle 3, 5 \rangle$ in i, j form

Example Write $2\mathbf{i} - 4\mathbf{j}$ in bracket vector form.

Lesson: The Dot Product

What is the dot product?

What is another name for the dot product?

Which is true: the dot product results in a vector

the dot product results in a real number

Example Find the dot products.

a. $\langle 3, 5 \rangle \cdot \langle 2, -1 \rangle$

b. $\langle 2, -3 \rangle \cdot \langle -4, -5 \rangle$

What are the properties of the dot product?

Examples Let $\mathbf{u} = \langle 1, 3 \rangle$, $\mathbf{v} = \langle 2, 4 \rangle$, $\mathbf{w} = \langle -1, 2 \rangle$. Evaluate the following.

a. $\mathbf{u} \cdot \mathbf{v} - \mathbf{u} \cdot \mathbf{w}$

b. $(3\mathbf{u}) \cdot \mathbf{w}$

What are the possible outcomes of the dot product?

What is the geometric interpretation of the dot product? Draw a sketch.

Example Find the angle between the vectors.

a. $\mathbf{u} = \langle 1, 3 \rangle, \mathbf{v} = \langle 2, 4 \rangle$

b. $\mathbf{u} = \langle 1, -2 \rangle, \mathbf{v} = \langle -1, -3 \rangle$

Explain how to interpret the dot product in the following scenarios ...

The Dot Product is ...	The Angle between the Vectors is ...
------------------------	--------------------------------------

Positive	
----------	--

Negative	
----------	--

Zero	
------	--

When the dot product is 0, what does this mean?

What's the better word for perpendicular? (fancier, makes you sound smart, you should definitely use it to impress people at parties):

Example Determine if the pair of vectors is orthogonal: $\mathbf{u} = \langle 1, 2 \rangle, \mathbf{v} = \langle -6, 3 \rangle$

Lesson: A Quick Review of the Basics of Solving Systems of Linear Equations

Systems of equations are 2 or more equations in 2 or more variables. They can look like:

$$\begin{cases} 2x + 3y = 6 \\ x - 3y = 3 \end{cases} \quad \text{or} \quad \begin{cases} 2x - y + z = 8 \\ 5x + 2y - z = 9 \\ x - 4y + z = 10 \end{cases} \quad \text{or} \quad \begin{cases} 3x + 2y - z = 7 \\ x - 4y + 2z = 8 \end{cases}$$

A solution of a system is an ordered pair that is true in each equation of the system. Some systems have no solutions, and some systems have infinite solutions.

In this lesson, we will briefly review the main techniques to solve systems. We will focus on systems that have a solution.

Example Use the substitution method to solve: $\begin{cases} 3x - y = -4 \\ x + 3y = 12 \end{cases}$

Example Use the elimination method to solve: $\begin{cases} 3x - y = -4 \\ x + 3y = 12 \end{cases}$

Example Solve:
$$\begin{cases} \frac{1}{2}x + \frac{1}{3}y = 4 \\ \frac{3x}{2} + \frac{3y}{2} = 15 \end{cases}$$

Example Solve:
$$\begin{cases} 2x + y + z = 9 \\ -x - y + z = 1 \\ 3x - y + z = 9 \end{cases}$$

Examples: Systems of 3 Linear Equations with 3 Variables

Solve:
$$\begin{cases} x + y + z = 2 \\ 2x + y - z = 5 \\ x - y + z = -2 \end{cases}$$

Solve:
$$\begin{cases} x + 2y + 3z = -11 \\ 3x - y + z = 0 \\ -2x + 3y - z = 4 \end{cases}$$

Lesson: Systems with No Solutions or Infinite Solutions

Example Solve:
$$\begin{cases} x + 4y - 3z = 2 \\ 2x - 5y + 2z = -8 \\ -3x - 12y + 9z = 7 \end{cases}$$

Example Solve:
$$\begin{cases} 7x + 2y = 6 \\ 14x + 4y = 12 \end{cases}$$

Example Solve:
$$\begin{cases} 5x - 4y + z = 9 \\ y + z = 15 \end{cases}$$

Example
$$\begin{cases} 3x - 2y + z = 15 \\ x + 4y - z = 11 \end{cases}$$

Example Solve:
$$\begin{cases} 2x + y - 3z = 0 \\ 4x + 2y - 6z = 0 \\ x - y + z = 0 \end{cases}$$

Lesson: Solving Systems of Equations by Graphing

We will now turn our attention to systems of equations.

System of Equations

A system of equations refers to 2 or more equations that share 2 or more variables.

Solution of a System of Equations

A solution of a system of equations is an ordered pair that is true in all equations.

Determine if the following pairs are solutions to the systems.

a. $(1,3)$

$$x - 2y = -5$$

$$2x - 3y = -4$$

b. $(-4,3)$

$$x + 3y = 5$$

$$2x + 3y = 1$$

There are several ways to solve systems. The first method we will focus on is how to solve by graphing.

Solving by Graphing


In this method, we graph both lines and determine the point of intersection if possible. We will examine all the possibilities.

Example

Graph each line and determine the point of intersection to find the solution.


$$y = 2x + 3$$

$$y = -\frac{1}{2}x - 2$$




Example

a. $y = -3x + 1$
 $y = 2x - 4$




b. $y - \frac{1}{2}x = -4$
 $y + 2x = 1$




There are two special situations that can occur with graphing.

Example

a. $y = -3x + 2$
 $9x + 3y = 6$



b. $y = -\frac{1}{2}x + 4$
 $2x + 4y = 12$



Lesson: Solving Systems of Equations by Substitution

In this lesson, we will focus on the second method for solving systems: substitution.

Show how to use substitution with this example:

$$2x + y = 6$$

$$3x + 5y = 9$$

Example Solve the system using substitution

$$x - 2y = 10$$

$$5x - 7y = 20$$

Below are 2 more complicated problems using substitution. Outline the steps shown in the video:

$$-2x + 4y = 4$$

$$-2x + 6y = 6$$

$$2x - 5y = -4$$

$$8x - 9y = 6$$

Now let's see what happens when we get a problem with no solution or infinite solutions.

a. $9y - 18x = 5$

$$2x - y = 3$$

b. $x + 2y = -3$

$$4x + 12 = -8y$$

Lesson: Solving Systems of Equations by Elimination

In this lesson, we will focus on the third method for solving systems: elimination.

Show how to use elimination with this example:

$$3x - y = 8$$

$$2x + y = -3$$

Example Use elimination to solve the system.

$$4x + 2y = -6$$

$$3x - 2y = 6$$

Now let's examine a more complicated case of elimination. Write down how to approach these problems:

a. $3x - 6y = 12$

$$5x - 2y = -4$$

b. $5x - 10y = 5$

$$2x - 3y = 6$$

Now let's investigate what happens in the situation where there is no solution or infinite solutions.

a. $5x - 2y = 1$
 $-10x + 4y = 3$

b. $3x + y = 5$
 $-9x - 3y = -15$

Lesson: Solving Systems of Equations that Require Clearing of Fractions

Examples

a.
$$\begin{aligned}\frac{1}{6}x - \frac{1}{3}y &= \frac{1}{2} \\ \frac{2}{5}x - \frac{3}{5}y &= -\frac{6}{5}\end{aligned}$$

b.
$$\begin{aligned}-\frac{1}{2}x - \frac{3}{4}y &= 1 \\ \frac{1}{3}x + \frac{3}{2}y &= -\frac{2}{3}\end{aligned}$$

c. $\frac{1}{2}x + \frac{3}{5}y = 6$

$$\frac{1}{4}x - \frac{1}{2}y = -9$$

Lesson: Solving Systems of Equations that Require Clearing of Decimals

Examples

a. $0.02x + 0.1y = 0.05$
 $4x + 20y = 10$

b. $0.1x - 0.05y = 0.2$
 $0.3x - 0.25y = 0.2$

c. $-0.4x + 1.3y = 1.$
 $1.6x - 0.9y = 6.9$

Lesson: Applications of Systems

The top of a rectangular desk is twice as long as it is wide. The perimeter of the desk is 180 inches. What are the dimensions of the desk?

Sarah bought 3 shirts and a scarf for \$53. Marie bought 2 shirts and 3 scarves for \$54. Find the cost of each item.

How many milliliters of a 5% acid solution and how many milliliters of a 17% acid solution must be mixed to obtain a 60 mL solution that is 15% acid?

Lesson: Systems of Equations with 3 Equations

Show how to solve the system:

$$x + 2y + 3z = -11$$

$$3x - y + z = 0$$

$$-2x + 3y - z = 4$$

Sometimes you will have missing variables. Show how to solve the following system that is missing some variables:

$$x - 2y = -3$$

$$x + 6z = 3$$

$$2y - 4z = 6$$

Now let's look at an example where there is no solution versus infinite solutions:

$$8x + 20y - 4z = 16$$

$$-2x - 5y + z = -4$$

$$10x + 25y - 5z = 20$$

$$x + 4y - 3z = 2$$

$$2x - 3y + 2z = -5$$

$$3x + 12y - 9z = 2$$

Examples: More Examples of Systems of 3 Equations

Example Solve the system:

$$x + 3y - z = -2$$

$$-x + 3y - 2z = 5$$

$$2x + 4y + 2z = 4$$

Example Solve the following system:

$$x - 2z = 2$$

$$-4y + 2z = -2$$

$$-x + 3y = -3$$

Example Solve the following system:

$$x - 4y + 2z = -7$$

$$3x - 8y + z = 7$$

$$6x - 12y + 3z = 12$$

Lesson: Using Matrices to Solve Systems of Linear Equations

What is a matrix?

Show how to setup the associated **augmented matrix** for the system: $\begin{cases} 3x + 4y = 4 \\ x - y = 13 \end{cases}$

What are the dimensions of this matrix?

In general, how do we state dimensions of a matrix (circle): rows x columns columns x rows

Example What is the dimension of

a.

$$\begin{bmatrix} 3 & 2 & 4 & 1 \\ 7 & 8 & 1 & 2 \\ 9 & 5 & -2 & -3 \end{bmatrix}$$

b.

$$\begin{bmatrix} 2 & 4 \\ 5 & 3 \\ 8 & 3 \\ -1 & -4 \end{bmatrix}$$

With matrices, there are 3 **row transformations** that we can make. We are not allowed to do anything else to a matrix. Write down these 3 transformations.

Now, recall, our goal in this lesson is to use a matrix to solve systems of linear equations. What do we call this specific solving process?

What do we call the “ideal form” of a matrix in its solved form?

The Gauss-Jordan method has a goal of transforming a matrix into reduced-row echelon form or diagonal form. Draw a few examples of how the FINAL matrix should look after we transform it:

When we use the Gauss-Jordan method, which direction do we want to work on the matrix?

What is the general strategy to remember with this method?

Example Outline the steps for the Gauss-Jordan method for the system: $\begin{cases} -2x + 4y = -4 \\ x - y = 4 \end{cases}$

Example Use Gauss-Jordan method for the system: $\begin{cases} 3x + 4y = 4 \\ x - y = 13 \end{cases}$

Lesson: Using The Gauss-Jordan Method for a System with 3 Equations

Example Use the Gauss-Jordan method for the system:
$$\begin{cases} x + y + z = 2 \\ 2x + y - z = 5 \\ x - y + z = -2 \end{cases}$$

Another page for you to show your work ...

Lesson: Using Matrices to Solve Systems of Linear Equations when There is No Solution or Infinite Solutions

Example Use the Gauss-Jordan method to solve:
$$\begin{cases} 3x + 5y - z = -2 \\ 4x - y + 2z = 1 \\ -6x - 10y + 2z = 0 \end{cases}$$

Example Use the Gauss-Jordan method to solve:
$$\begin{cases} 3x + 5y = -2 \\ 9x + 15y = -6 \end{cases}$$

Example Use the Gauss-Jordan method to solve: $\begin{cases} x - 2y + 3z = 6 \\ 2x - y + 2z = 5 \end{cases}$

Example Use the Gauss-Jordan method to solve:
$$\begin{cases} 5x - 4y + z = 0 \\ x + y = 0 \\ -10x + 8y - 2z = 0 \end{cases}$$

Examples: Using Matrices to Solve Systems of Linear Equations (any type of solutions)

Use the Gauss-Jordan method to solve the following.

a.
$$\begin{cases} x + 4y - z = 6 \\ 2x - y + z = 3 \\ 3x + 2y + 3z = 9 \end{cases}$$

b.
$$\begin{cases} 2x + y - 3z = 0 \\ 4x + 2y - 6z = 0 \\ x - y + z = 0 \end{cases}$$

c.
$$\begin{cases} 3x - 2y + z = 15 \\ x + 4y - z = 11 \end{cases}$$

Lesson: Nonlinear Systems of Equations

Nonlinear systems of equations involve at least one equation that is not linear. Some examples look like this:

$$\begin{cases} x^2 - y = 8 \\ x + y = -4 \end{cases} \quad \text{or} \quad \begin{cases} x^2 + y^2 = 4 \\ |x| + y = 2 \end{cases}$$

Similar to systems of linear equations, we can still use substitution or elimination to solve the system. However, the best method to use tends to depend on the type of system you are trying to solve. Additionally, you might have to use both methods.

What is one potential issue that can happen with solving systems of nonlinear equations and what should you do about this?

When one equation is linear and the other is not, it is usually best to use the substitution method.

Example Solve the system: $\begin{cases} x^2 + y = 8 \\ x - y = -2 \end{cases}$

When both variables in both equations are squared, it's usually best to use elimination.

Example Solve the system: $\begin{cases} x^2 + y^2 = 25 \\ x^2 + 2y^2 = 9 \end{cases}$

Sometimes you will have to use both the substitution and elimination methods.

Example Solve the system: $\begin{cases} x^2 - xy + y^2 = 5 \\ 2x^2 + xy - y^2 = 10 \end{cases}$

Now let's look at a few other examples to get a sense of how to use these methods!

Example Solve the system: $\begin{cases} x^2 + y^2 = 9 \\ -|x| + y = 0 \end{cases}$

Example Solve the system: $\begin{cases} 3x^2 + y^2 = 3 \\ 4x^2 + 5y^2 = 26 \end{cases}$

Lesson: Properties of Matrices: Equality, Addition, Subtraction, and Scalar Multiplication

When are 2 matrices equal?

Example What values of x , y , z , and w will make the matrices equal?

$$\begin{bmatrix} 2+x & 3-w \\ 7 & 3 \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 2-y & 2+z \end{bmatrix}$$

When can you add 2 matrices?

What is the definition of matrix subtraction?

What is scalar multiplication with a matrix?

List the properties of scalar multiplication for matrices:

Example Let $A = \begin{bmatrix} 3 & 2 \\ 1 & 8 \end{bmatrix}$, $B = \begin{bmatrix} 7 & 2 & 1 \\ 0 & 4 & 1 \end{bmatrix}$, $C = \begin{bmatrix} 0 & 7 \\ 1 & 3 \end{bmatrix}$, $D = \begin{bmatrix} 2 & 3 \\ 1 & 2 \\ 1 & 0 \end{bmatrix}$, $E = \begin{bmatrix} 7 & 2 & 1 \\ 1 & 0 & 0 \end{bmatrix}$. Complete the following operations, if possible.

a. $A + C$

b. $B - D$

c. $B - E$

d. $2B + E$

e. $2A - 3C$

Lesson: Matrix Multiplication

First, let's review dimension. What is the dimension of the following matrices?

$$A = \begin{bmatrix} 3 & 2 \\ 1 & 8 \end{bmatrix}$$

$$B = \begin{bmatrix} 7 & 2 & 1 \\ 0 & 4 & 1 \end{bmatrix}$$

$$D = \begin{bmatrix} 2 & 3 \\ 1 & 2 \\ 1 & 0 \end{bmatrix}$$

To understand how to multiply matrices, let's use the following 2 examples:

a. $\begin{bmatrix} 2 & 7 \\ -4 & 3 \end{bmatrix} \begin{bmatrix} 5 & -1 \\ 3 & 2 \end{bmatrix}$

b. $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 1 & 0 \\ 7 & 2 \end{bmatrix}$

Example Let $A = \begin{bmatrix} 3 & 2 \\ 1 & 8 \end{bmatrix}$, $B = \begin{bmatrix} 7 & 2 & 1 \\ 0 & 4 & 1 \end{bmatrix}$, $C = \begin{bmatrix} 0 & 7 \\ 1 & 3 \end{bmatrix}$, $D = \begin{bmatrix} 2 & 3 \\ 1 & 2 \\ 1 & 0 \end{bmatrix}$, $E = \begin{bmatrix} 7 & 2 & 1 \\ 1 & 0 & 0 \end{bmatrix}$. Complete the following operations, if possible.

a. AC

b. CA

Using the above example, what can we say about matrix multiplication and commutativity?

List the properties of matrix multiplication:

Example Let $A = \begin{bmatrix} 3 & 2 \\ 1 & 8 \end{bmatrix}$, $B = \begin{bmatrix} 7 & 2 & 1 \\ 0 & 4 & 1 \end{bmatrix}$, $C = \begin{bmatrix} 0 & 7 \\ 1 & 3 \end{bmatrix}$, $D = \begin{bmatrix} 2 & 3 \\ 1 & 2 \\ 1 & 0 \end{bmatrix}$, $E = \begin{bmatrix} 7 & 2 & 1 \\ 1 & 0 & 0 \end{bmatrix}$. Complete the following operations, if possible.

a. BC

b. ED

c. CB

Lesson: Inverse Matrices

What is the 2×2 identity matrix?

What is the 3×3 identity matrix?

How do we describe the general $n \times n$ identity matrix?

Example Multiply $A = \begin{bmatrix} 3 & 2 & 1 \\ 8 & 1 & 7 \\ 3 & 1 & 0 \end{bmatrix}$ by the identity matrix as follows:

a. AI

b. IA

What can we generalize about the identity matrix?

How do we describe a multiplicative inverse of a real number?

Some matrices have inverses. For a matrix A , what notation denotes its inverse?

What is the property of inverse matrices?

What are the steps for finding an inverse matrix?

Example Find the inverses of the following matrices. Prove your answer by checking it: $\begin{bmatrix} 6 & -2 \\ 2 & 4 \end{bmatrix}$

Example Find the inverses of the following matrices. Prove your answer by checking it: $\begin{bmatrix} 3 & 1 & 2 \\ 1 & -1 & 1 \\ 2 & 1 & -1 \end{bmatrix}$

Example Find the inverses of the following matrices. Prove your answer by checking it: $\begin{bmatrix} 3 & 2 \\ 6 & 4 \end{bmatrix}$

Lesson: Using Inverse Matrices to Solve Systems

We've previously talked about using matrices to solve systems. We will look at another way to do that in this lesson. Show another way to set up the system using matrices (as well as the relevant notation):

$$\begin{cases} x + y + z = 6 \\ 2x + 3y - z = 7 \\ 3x - y - z = 6 \end{cases}$$

Now, using our setup above, explain the reasoning for how inverse can help solve a system:

Example Use inverse matrices to solve: $\begin{cases} -x + y = 1 \\ 2x - y = 1 \end{cases}$

Example Solve the system: $\begin{cases} x + y + z = 6 \\ 2x + 3y - z = 7 \\ 3x - y - z = 6 \end{cases}$ using inverse matrices.

Lesson: How to Calculate the Determinate of a Matrix

Every $n \times n$ matrix has a real number associated with it called the **determinant**. What is the notation to represent this?

Example Show, in detail, how to calculate the determinants of the following matrices.

a. $A = \begin{bmatrix} 2 & 4 \\ -5 & 7 \end{bmatrix}$

b. $B = \begin{bmatrix} -5 & 1 \\ -3 & 4 \end{bmatrix}$

Example Show, with illustrations, how to calculate the determinant of $A = \begin{bmatrix} 1 & 2 & 3 \\ -2 & 1 & -3 \\ 3 & 4 & -2 \end{bmatrix}$

The 2x2 “sub” matrices that we calculate are called minors. The symbol M_{ij} represents the minor that results when row i and column j are eliminated. Using the matrix above, describe the following minors.

$$M_{11} =$$

$$M_{32} =$$

The number that we multiply the minor against is called the **cofactor**. Draw an illustration to show whether a cofactor will be positive or negative:

Example Using the matrix below, describe the minor associated with each cofactor AND determine whether the cofactor is positive or negative. You do not need to calculate the determinant.

$$A = \begin{bmatrix} 2 & 4 & 6 \\ 1 & 3 & 5 \\ -3 & -5 & -1 \end{bmatrix}$$

a. 2

b. 3

c. -3

Example Show how to efficiently calculate the determinants of the following matrices. EXPLAIN YOUR REASONING FOR WHY YOU DID IT THE WAY THAT YOU DID.

a. $A = \begin{bmatrix} 2 & 7 & 8 \\ -1 & 0 & 1 \\ 3 & 5 & 1 \end{bmatrix}$

b. $B = \begin{bmatrix} 9 & -3 & 1 \\ 2 & 3 & 0 \\ 5 & 2 & 0 \end{bmatrix}$

Lesson: Determinant Theorems

Theorem 1: If every element (or column) of a matrix is 0, then the determinant will be 0.

Example: find the determinant of $A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 4 & 2 & 8 & 9 \\ 3 & 2 & 8 & 4 \\ 9 & 3 & 1 & 7 \end{bmatrix}$

Theorem 2: If the rows of a matrix A are the corresponding columns of a matrix B, then the determinants are equal.

Example: Calculate both determinants: $A = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 1 \\ 3 & 2 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 3 & 1 & 0 \end{bmatrix}$

Theorem 3: If any 2 rows (or columns) of a matrix are interchanged, then the determinants are negatives of one another.

Example Find the determinants of: $A = \begin{bmatrix} 2 & 4 \\ 5 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 5 & 6 \\ 2 & 4 \end{bmatrix}$

Theorem 4: Suppose matrix B is formed by multiplying every element in a row or columns in a matrix A by a real number k. Then the determinant is $|B| = k|A|$.

Example: Find the determinants of: $A = \begin{bmatrix} 2 & 1 \\ 3 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 8 & 4 \\ 3 & 6 \end{bmatrix}$

Theorem 5: If two rows (or columns) of a matrix are identical, then the determinant will be 0.

Example Find the determinant of: $A = \begin{bmatrix} 4 & 4 \\ 4 & 4 \end{bmatrix}$

Theorem 6: Changing a row (or column) of a matrix by adding it to a constant times another row (or column) does not change the determinant of the matrix.

Example Find the determinant of $A = \begin{bmatrix} 1 & 2 \\ 5 & 8 \end{bmatrix}$. Then create a matrix B by taking adding $3 * (\text{row } 3)$ to row 1. Find the new determinant.

Theorem 7: If a matrix is in triangular form (all 0s either above or below the diagonal), then the determinant is found by multiplying the elements down the diagonal.

Example: Find the determinant of $A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 3 & 5 & 0 & 0 \\ 2 & 1 & 2 & 0 \\ 8 & 7 & 3 & 1 \end{bmatrix}$

Examples Use the determinant theorems to find the following determinants.

a. $\begin{vmatrix} -4 & 1 & 4 \\ 2 & 0 & 1 \\ 0 & 2 & 4 \end{vmatrix}$

b. $\begin{vmatrix} 1 & 0 & 8 \\ 4 & 0 & 2 \\ 0 & 0 & 8 \end{vmatrix}$

Lesson: Ellipses as Conic Sections

Give the geometric definition of an ellipse. Draw an illustration to accompany this.

What are the standard forms of an ellipse? Describe the parts and draw VERY CLEAR illustrations.

What is important to note with this equation?

Example Graph each ellipse. Give the domain, range, center, vertices, endpoints of the minor axis, and focus.

a. $\frac{x^2}{9} + \frac{y^2}{25} = 1$

b. $9x^2 + y^2 = 81$

Example Graph each ellipse. Give the domain, range, center, vertices, endpoints of the minor axis, and focus.

c. $\frac{(x-1)^2}{9} + \frac{(y+2)^2}{4} = 1$

Lesson: Writing Equations of Ellipses

Example Write the equation of an ellipse with the following characteristics.

a. major axis 6, focus at $(0, 2)$ and $(0, -2)$

b. center at $(-2, 7)$, major axis vertical with length 10, and $c=2$

Lesson: Eccentricity with Ellipses

To discuss the concept of eccentricity, we need to discuss what the formal mathematical definition of conic sections is.

Define a conic:

The intuitive idea behind eccentricity can change slightly with each conic section, but it is generally a ratio of different measures within the conic section.

What does the eccentricity of an ellipse tell us?

How do we represent eccentricity?

How do you calculate the eccentricity of an ellipse?

How does the eccentricity of a parabola compare to the eccentricity of an ellipse? Summarize the explanation from the video.

Examples Find the eccentricity of each ellipse.

a. $\frac{x^2}{9} + \frac{y^2}{4} = 1$

b. $x^2 + 25y^2 = 25$

Example Find the equation of an ellipse with $e = \frac{3}{4}$ with foci at $(-2,0), (2,0)$

Examples: Using Complete the Square to Write an Ellipse Equation in Standard Form

Rewrite the following equations in standard form.

a. $x^2 + 12y^2 - 14x - 48y = -85$

b. $9x^2 + 36x + 16y^2 - 160y + 292 = 0$

Lesson: Hyperbolas as Conic Sections

What is the geometric definition of a hyperbola? Draw an illustration with your definition.

State the equations for the standard form of the hyperbola. Draw illustrations with it.

Examples Graph each hyperbola. Give the domain, range, center, vertices, foci, and equations of the asymptotes.

a. $\frac{x^2}{9} - \frac{y^2}{16} = 1$

b. $9x^2 - 25y^2 = 225$

Examples Graph each hyperbola. Give the domain, range, center, vertices, foci, and equations of the asymptotes.

c. $\frac{(y-1)^2}{9} - \frac{(x+3)^2}{4} = 1$

Lesson: Writing Equations of Hyperbolas

Examples Write the equation of the hyperbola with the following characteristics.

a. vertices at $(0, 6)$ and $(0, -6)$ with asymptotes $y = \pm \frac{1}{2}x$

b. foci at $(-3\sqrt{5}, 0)$, $(3\sqrt{5}, 0)$ with asymptotes $y = \pm 2x$

Examples Write the equation of the hyperbola with the following characteristics.

c. center at $(1, -2)$ and focus at $(-2, -2)$ and vertex at $(-1, -2)$

Lesson: Eccentricity of Hyperbolas

How do you calculate the eccentricity of a hyperbola?

What can we say about the eccentricity of the hyperbola? What does it mean?

Example Find the eccentricity of the hyperbolas.

a. $\frac{x^2}{36} - \frac{y^2}{9} = 1$

b. $8y^2 - 16x^2 = 16$

Example Find the equation of a hyperbola with $e = \frac{25}{9}$ and foci at (9, -1) and (-11, -1)

Examples: Writing the Standard Form of a Hyperbola

Write each hyperbola in standard form.

a. $x^2 - 6x - y^2 + 4y = 1$

b. $y^2 - 14y - x^2 + 14x = 16$

Lesson: An Intro to Conic Sections

What are the 4 conic sections we will discuss? Draw an illustration with it.

Lesson: Parabolas as Conic Sections

What is the previous equation we used for parabolas opening upwards or downwards? Draw an illustration to go with this.

How can we represent parabolas in similar notation for parabolas that open sideways?

Examples Give the domain and range of the following parabolas, then graph the parabola.

a. $x + 1 = (y + 1)^2$

b. $-x = 3y^2 + 6y + 2$

When dealing with conic sections, we shift how we think about things.

What is the geometric definition of a parabola? Draw an illustration to accompany this definition.

What equations do we use that align with the geometric definition?

Examples Give the focus, directrix, vertex, and axis of symmetry for each parabola. Then use this information to graph the parabola.

a. $x^2 = 4y$

b. $(y - 2)^2 = 24(x - 3)$

Lesson: Writing Equations of Parabolas

Example Write the equation of a parabola with the following characteristics.

a. focus $\left(0, -\frac{1}{2}\right)$ with the vertex at the origin

b. vertex $(4, 3)$ and focus is $(4, 5)$

c. vertex $(1, 2)$, directrix $x=4$

Lesson: Sequences

Define a sequence.

Examples Write the first five terms of each sequence.

a. $a_n = \frac{1}{n}$

b. $a_n = \frac{n+1}{n+3}$

c. $a_n = (-1)^{n+1} * (2n + 1)$

What does it mean for a sequence to converge?

What does it mean for a sequence to diverge?

What is a recursively defined sequence?

Examples Find the first four terms in the sequence.

a. $a_1 = 3; a_n = 2a_{n-1} - 4$

b. $a_1 = 2; a_n = a_{n-1}^2 + n - 2$

Lesson: Series

What is the summation sign and what does it mean? What do we call the “ i ” in this formula?

Define a finite series.

Define an infinite series.

Examples Evaluate the following sums.

a. $\sum_{i=1}^4 (3^i + 1)$

b. $\sum_{j=3}^6 a_j$

c. $\sum_{k=1}^4 (4x_k - 1)$ if $x_1 = 2, x_2 = 3, x_3 = 5, x_4 = 8$

d. $\sum_{i=1}^3 f(x_i) \Delta x$, if $f(x) = 3x, x_1 = 1, x_2 = 4, x_3 = 7, \Delta x = 3$

What are the summation properties?

What are the summation rules for summing up terms?

Examples Use the summation properties and rules to evaluate each series.

a. $\sum_{i=1}^{50} 3$

b. $\sum_{k=1}^{25} 3k$

c. $\sum_{j=1}^{33} (4j^2 + 1)$

Lesson: Arithmetic Sequences

What is an arithmetic sequence?

Give some examples of an arithmetic sequence.

Example Determine the common difference, d , for the arithmetic sequence:

$$-7, -5, -3, -1, \dots$$

Example Find the first five terms of each arithmetic sequence.

a. The first term is 3, and the common difference is -2.

b. $a_1 = -5, d = 7$

How do we define the general n th term of an arithmetic sequence?

Example Determine a_{11} and a_n for the arithmetic sequence $-2, 2, 6, 10, \dots$

Example Determine a_n and a_{20} for the arithmetic sequence having $a_3 = 9$, and $a_4 = 16$.

Example An arithmetic sequence has $a_7 = -15$ and $a_{14} = -35$. Determine a_1 .

Lesson: Arithmetic Series

When we sum up the terms of an arithmetic sequence, how can we define that sum? (sum of an arithmetic sequence)

Example Consider the arithmetic sequence $-7, -3, 1, 5, \dots$

a. Evaluate S_{10}

b. Evaluate the sum of the first 50 positive integers.

Example The sum of the first 15 terms of an arithmetic sequence is 345. If $a_{15} = 65$, find a_1 and d .

Examples Evaluate each sum.

a. $\sum_{i=1}^{19} (-3 - 5i)$

b. $\sum_{k=4}^{15} (2 + 4k)$

Lesson: Geometric Sequences and Series

Define a geometric sequence:

Example Determine a_6 and a_n for the geometric sequence 4, 12, 36, 108,

Example Determine r and a_1 for the geometric sequence with $a_2 = -18$ and $a_5 = 486$

How do we define the sum of the first n terms of a geometric sequence?

Example Evaluate: $\sum_{i=1}^8 3 * 2^i$

If we start with 2 and use a common ratio of $\frac{1}{2}$, what eventually happens as the series continues infinitely? Summarize the explanation below.

Summarize the Sum of the Terms of an Infinite Geometric Sequence

Examples Evaluate the following infinite sums.

a. The first term is 1 and the common ratio is $\frac{1}{3}$

b. $\sum_{i=1}^{\infty} \left(-\frac{1}{2}\right) \left(-\frac{3}{4}\right)^{i-1}$

c. $\sum_{i=1}^{\infty} \left(\frac{3}{7}\right)^i$

Lesson: The Binomial Theorem

Below we can observe the pattern for binomial expansion:

$$(x + y)^0 = 1$$

$$(x + y)^1 = x + y$$

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

$$(x + y)^5 = x^5 + 5x^4y + 10x^3y^2 + 10x^2y^3 + 5xy^4 + y^5$$

Focusing on the VARIABLES x and y only, how can we generalize the exponents for the variables?

Pascal's Triangle can explain how to find the coefficient. Draw out Pascal's triangle below.

Having to draw out this pattern would be very difficult for larger expansions, so we will explore a few ways to simplify how to expand this pattern.

First, we need to understand factorials.

n-factorial: for any positive integer n ,

$$n! = n(n - 1)(n - 2) \dots (3)(2)(1)$$

By definition: $0! = 1$

Examples Evaluate:

a. $5!$

b. $8!$

Now that we understand factorials, we can discuss the formula to find binomial coefficients.

For nonnegative integers n and r , $r \leq n$, the binomial coefficient is defined as follows:

Examples Evaluate each binomial coefficient:

a. $\binom{5}{2}$

b. $\binom{6}{0}$

We can summarize how to expand $(x + y)^n$ as follows:

1. There are $n+1$ terms in the expansion
2. The first and last terms are x^n and y^n respectively
3. In each succeeding term, the exponent on x decreases by 1 and the exponent on y increase by 1
4. The sum of the exponents on x and y in any term is n .
5. The coefficient of the term with $x^r y^{n-r}$ or $x^{n-r} y^r$ is $\binom{n}{r}$

These observations are captured in the binomial theorem.

State **The Binomial Theorem**:

Example Find the first, last, and seventh term of the binomial expansion: $(3x + 2y)^{10}$

Example Use the binomial theorem to expand: $(a - b)^5$

Not included (no notes file):

004 (Examples of the Greatest Common Factor)

005 (Examples of Factoring Basic Trinomials via Trial and Error)

006 (Examples of Grouping)

007 (Examples of the Sum and Difference of Cubes)

008 (Examples of Factoring Harder Trinomials)

013 (Examples of Simplifying Rational Expressions)

014 (Examples of How to Use the Multiplicative Property of -1 to Simplify Rational Expressions)

015 (Examples of Finding Domains of Rational Expressions)

018 (Examples of Multiplying and Dividing Rational Expressions)

022 (Examples of Finding Common Denominators for Rational Expressions)

037 (Examples of Solving Absolute Value Equations)

042 (Examples on Absolute Value Inequalities)

878 (Examples: Putting Everything Together and Another Explanation of Special Cases, Slower-Paced)